

Evaluation of the Role of Serum Malondialdehyde in the Pathogenesis of Diabetic Retinopathy

DOI: <https://doi.org/10.32007/jfacmedbagdad.6431957>

Zena M. Hassan *

BSc, MSc

Rana A. Hamdi *

BSc, PhD

Ebtehal N. Al Bassam **

MBChB,DO



This work is licensed under a [Creative Commons Attribution-NonCommercial 4.0 International License](https://creativecommons.org/licenses/by-nc/4.0/).

Abstract:

Background: The most typical consequence of diabetes mellitus is diabetic retinopathy. An important part of the etiology of diabetes and diabetic retinopathy is played by oxidative stress. Malondialdehyde is a dialdehyde that is moderately toxic and is frequently used as a marker for oxidative stress as well as lipid peroxidation.

Objectives: To assess the serum malondialdehyde levels in diabetic patients with and those without retinopathy and to compare these levels to controls. In additions to, investigate the relationship between serum malondialdehyde level and long glycemic control, the glycated hemoglobin (HbA1c).

Patients and Methods: This case-control study included (120) individuals from 40 to 70 years of age. They were divided into three groups: Group 1: 40 type 2diabetic patients with retinopathy, Group 2 :40 type 2 diabetic patients without retinopathy, and Group 3 40 controls .The biochemical testes included fasting blood glucose measured by Cobas c 311 systems, glycated hemoglobinHbA1c measured by Cobas c 111 systems, and serum malondialdehyde (MDA) measured by enzyme linked immunosorbent assay (ELISA).

Results: A higer mean value of (MDA) was found serum diabetic patients with and without retinopathy as compared to control ($p=0.0001$). As well as a significantly higher mean value of serum (MDA) in diabetic patients with retinopathy as compared to those without ($p=0.0001$).A significant positive correlation was found between serum(MDA) and HbA1C in diabetic patient with retinopathy group ($r=0.931, p=0.0001$).

Conclusion: Higher serum levels of malondialdehyde is an indicator of increased lipid peroxidation that may be involved in pathogeneses of retinopathy in uncontrolled type 2 diabetic patients.

Keywords: Oxidative stress, diabetic retinopathy, malondialdehyde

Introduction:

Diabetes mellitus (DM) is a chronic condition that is defined by an elevated blood sugar level. It happens either when the body does not produce enough insulin or when the cells do not respond to the insulin that is produced (1,2,3). DM is associated with many complications (4). Retinopathy is one of DM's long-term consequences, which may result in vision loss, Nephropathy, atherosclerotic cardiovascular disease, peripheral neuropathy, arterial disease, as well as cerebral vascular disease are also possible complications of DM (5). Diabetic retinopathy (DR) a known cause of blindness in people in the working age, and is one of the most important pathologic vascular effects of diabetes (6). The pathophysiology of retinal microvascular injury is heavily influenced by hyperglycemia. (7).

as it accelerates the development of oxidative stress and increases plasma free radical concentration is hyperglycemia (8). Through the production of free radicals, particularly reactive oxygen species (ROS) (4). Vascular leakage, vascular malfunction, and pathological angiogenesis are all indicators of (DR) and are influenced by the overproduction of ROS (9). The ability of biological systems to detoxify (ROS) is out of balance during oxidative stress, which causes these substances to accumulate (10). Oxidative stress, and ROS, can damage DNA, proteins, and lipids, causing adjustments to DNA and proteins as well as lipid peroxidation. (MDA) is a by-product of polyunsaturated fatty acid peroxidation (11). And is an extremely dangerous chemical that develops from lipid peroxidation as a result of free radical damage (9). MDA is associated with oxidative stress and poor antioxidant defense, which promotes the progression of DR (12). Serum MDA level is a sensitive marker of lipid peroxidation that is a useful measure of oxidative stress status (11). Since hyperglycemia is the primary

*Dept of Biochemistry, College of Medicine, University of Baghdad, Email: Mzena4244@gmail.com

rana.chemist2006@yahoo.com

**Diabetes of specialized center of endocrinology, Email: ebtahalbassam601@gmail.com

cause of lipid peroxidation, the rise in serum MDA shows that this rate of MDA has increased (13).

Cases and Methods:

This case – control study included (120) individuals from 40 to 70 years of age. Participants were old diabetic but new retinopathy included all patients coming to the center of clients of the Diabetic Control Clinic and Diabetes/ Specialized Center for Endocrinology and Ibn- Al Haitham Teaching Hospital in Rusafa city in Baghdad .November 2021 to January 2022 Informed consent was obtained from each participant. The study was approved by the Scientific Committee of the College of Medicine/ University of Baghdad. Individuals were divided into three groups: 40 type 2 diabetics with retinopathy in Group 1, 40 type 2 diabetics with no retinopathy in group2. And 40 controls group three.

Patients with a history of severe eye illness or retinal detachment, type1DM, end-stage renal disease, malignancies, end-stage cardiac disease, immunosuppressive drug usage in the past or present, or IV drug steroid users were excluded from the study. The diagnosis of DM was based on the history of type2 DM, fasting serum glucose (FSB > 126) and glycated hemoglobin (HbA1C > 6.5%) according to the WHO criteria. The diagnosis of DR was made by after an ophthalmologist through history taking clinical and ophthalmological examination which. Included examination of the fundus by a slit lamp bio-microscope and indirect ophthalmoscope with 90-D lens, fundus color photograph centered on the macula, and optical coherence tomography (OCT).

Blood tests included glycated hemoglobin (HbA1C) measured by autoanalyzer (cobs) C111, fasting serum glucose (FSG) by autoanalyzer (cobs) C311 and serum MDA levels by an enzyme -linked immunoassay (ELISA).

Statistical analysis:

Data analysis was done using SPSS-27(Statistical Packages for Social Sciences- version 27) Simple statistics such as mean, standard deviation, and range were used to describe the groups. When comparing

two independent means, the Students’t-test was used, and when comparing more than two means, the ANOVA test was used. A P value of less than 0.05 was considered as statistically significant

Results:

The mean values of FBS and HbA1C in the two diabetic patients groups (with and without retinopathy) were significantly higher $p= 0.0001$ than the controls figures (1 and 2).




Figure (1): Mean value of FBS in all studied groups.




Figure (2): Mean value of HbA1c in every studied group.

The mean value of serum malondialdehyde in diabetic patients with and without retinopathy was significantly higher than the controls ($p=0.0001$). The mean value of serum malondialdehyde in diabetic patients with retinopathy significantly higher than those without retinopathy ($p=0.0001$ to those without retinopathy ($p=0.0001$), in table (1).

Table (1): Mean value of serum Malondialdehyde level in the studied groups.

	T2DM Retinopathy	T2DM	Control	P value
Mean±SD	27.94±4.45	15.84±2.48	5.2±1.0	
(Range)	(20.2 -35.7)	(10.34-18.9)	(2.76-6.91)	4
Malondialdehyde (MDA) (nmol/ml)	ANOVA	DR x C	DM x C	DR x DM
	0.0001^	0.0001#	0.0001#	0.0001#

Significant Positive correlation was found between serum MDA and HbA1C in diabetic patient with retinopathy group ($r=0.931, p=0.0001$), in Figure 3.




Figure (3) The correlation between MDA and HbA1C ($r=0.931, p=0.001$) in the diabetic retinopathy group

Discussion:

Hyperglycemia causes oxidative stress that is related to diabetes pathophysiology. Strong oxidative stress may reason harm to cells leading to microvascular issues that involve diabetes retinopathy (14). MDA is a three-carbon dialdehyde with a high reactivity that is created as a by-product of lipid peroxidation (9). It is a marker system that can be utilized to detect complications associated with diabetes (15). The result of this study revealed a higher level of serum MDA in diabetic both with and without retinopathy. This finding is in agreement with Dave *et al.* who showed the MDA levels higher level in patients with DM than in controls. A higher level of oxidants is found in DM, as a result (16). Moreover, a recent study revealed higher level of MDA in DR in comparison to both DM without retinal damage and control (suggesting that the high level of MDA in diabetic groups result from oxidative stress-induced lipid peroxidation (9)). Oxidative stress is important for the development of diabetes' pathology (20). Oxidative stress negatively affects the insulin's functions via numerous pathways interacting and producing ROS. These might weaken the pancreatic islet-cells, which would cause them to release less insulin. Moreover, free radical production via non-enzymatic protein glycation, enhanced lipid peroxidation and glucose oxidation results in cellular injury machinery, alterations to the cell membrane, enzyme damage increased insulin resistance and diabetes risk (17,18). The current study found positive correlation between MDA and HbA1c which is in line with recent research that found a connection between MDA and HbA1c. (9, 19). Uncontrolled diabetes is related to a greater MDA as a lipid peroxidation, indicator perhaps as a result of chronically elevated blood sugar followed by increased

oxidative stress. Shawki *et al.* found that MDA is a lipid peroxidation marker that interacts with the phospholipids in cell membranes (9).

Conclusions:

Higher serum levels of malondialdehyde is an indicator of increased lipid peroxidation that may be involved in the pathogenesis of retinopathy in uncontrolled type 2 diabetic patients.

Authors' Contributions:

Idea, design, and critical revision of the study: Zena Mohammed Hassan, dr Rana Ali Hamdi. Data gathering, analysis, and interpretation are all steps in the writing of a manuscript. : Zena Mohammed Hassan, dr Rana Ali Hamdi and diagnosis and Samples were provided by: dr Ebtehal Nouri Al Bassam

References:

- 1- Alwan FJ, Saleh AZ, Al-Najjar HY. Echocardiographic assessment of the effect of type (2) Diabetes mellitus on cardiac performance. *Journal of the Faculty of Medicine Baghdad* 2012; 54(3):252-255.
- 2- ADA. *Diagnosis and classification of diabetes mellitus. Diabetes Care* 2010; 33(Suppl1):S62-69..
- 3- Gouliopoulos NS, Kalogeropoulos C, Lavaris A, Rouvas A, Asproudis I, Garmpi A, *et al.* Association of serum inflammatory markers and diabetic retinopathy: a review of literature. *Eur Rev Med Pharmacol Sci* 2018; 22(21): 7113-7128.
- 4- Altoum AEA, Osman AL, Babker AM. Correlation of oxidative stress markers malondialdehyde (MDA), antioxidant vitamins A, E, and C with glycated hemoglobin (HbA1C) levels in Type 2 diabetes mellitus. *Asian J Pharm Clin Res.* 2018; 11(5):281-283.
- 5 -ADA. *Standards of medical care in diabetes 2018: Microvascular complication and foot care. Diabetes Care;* 2018; 41(Suppl. 1):S109-S111.
- 6- Elbana KA, Salem HM, Fattah NRA, Etman E. Serum pentraxin 3 level as a recent biomarker of diabetic retinopathy in Egyptian patients with diabetes. *Diabetes & Metabolic Syndrome: Clinical Research & Reviews.* 2019; 13(4):2361-2364
- 7- Ahsan H. Diabetic retinopathy-biomolecules and multiple pathophysiologies. *Diabetes Metab Syndr.* 2015;9(1):51-4.
- 8 - Mohammed HA, Al-Shamma GA, Hashim HM. Beta-carotene, glycemic control and dyslipidemia in type 2 diabetes mellitus. *Journal of the Faculty of Medicine Baghdad.* 2006; 48(4):435-441
- 9- Shawki HA, Elzehery R, Shahin M, Abo-Hashem EM, Youssef MM. Evaluation of some oxidative markers in diabetes and diabetic retinopathy. *Diabetology international.* 2021; 12(1):108-117..
- 10- Augustine J, Troendle EP, Barabas P, McAleese CA, Friedel T, Stitt AW, *et al.* The role of lipoxidation in

the pathogenesis of diabetic retinopathy. *Frontiers in endocrinology*. 2021; 11(article 621938):1-24.

11- Pan HZ, Zhang H, Chang D, Li H, Sui H. The change of oxidative stress products in diabetes mellitus and diabetic retinopathy. *British Journal of Ophthalmology*. 2008; 92(4):548-551

12- Khalaf MR, Ahmeid MS, Abdulqader OJ, Jabar HA. Evaluation of the Role of Malondialdehyde in Occurrence and Development of Diabetic Retinopathy Patients. *Medico-Legal Update*. 2021; 21(1):1400-1408.

13- El-Yassin HD, Ghassan AS. Oxidative DNA damage and lipid peroxidation as markers of oxidative stress type 2 diabetic patients. *Journal of the Faculty of Medicine Baghdad*. 2008; 50(1):110-119.

14- Abbasa SAN, Razaa ST, Mird SS, et al. Association of variants rs7903146 and rs290487 of TCF7L2 gene with diabetic nephropathy and co-morbidities (hypertension and dyslipidemia) in type 2 diabetes mellitus. *Meta Gene*. 2019;20(ID 00561):1-6.

15- Cecilia OM, José Alberto CG, José NP, Ernesto Germán CM, Ana Karen LC, Luis Miguel RP, et al. Oxidative stress as the main target in diabetic

retinopathy pathophysiology. *Journal of diabetes research*, 2019; 2019(ID 8562408):1-21.

16- Dave A, Kalra P, Gowda BH, Krishnaswamy M. Association of bilirubin and malondialdehyde levels with retinopathy in type 2 diabetes mellitus. *Indian J EndocrinolMetab*. 2015; 19(3):373-7.

17- Asmat U, Abad K, Ismail K. Diabetes mellitus and oxidative stress—a concise review. *Saudi Pharm J*. 2016; 24(5):547–53.

18- Dos Santos JM, Tewari S, Mendes RH. The role of oxidative stress in the development of diabetes mellitus and its complications. *J Diabetes Res*. 2019; 2019(4189813):1-3.

19- Zarei M, Farahnak Z, Hosseinzadeh-Attar MJ, et al. Lipid peroxidation and antioxidant enzymes activity in controlled and uncontrolled Type 2 diabetic patients. *ARYA Atheroscler*. 2016; 12(3):118-123.

20- Korany MA, Sonbol A, Elgouhary SM. Omentin-1 and diabetic retinopathy in type 2 diabetic patients. *Alexandria J Med*. 2018; 54(4):323–326.

How to cite this Article:

Hassan ZM, Hamdi RA, Al Bassam EN. Evaluation of the Role of Serum Malondialdehyde in the Pathogenesis of Diabetic Retinopathy. *JFacMedBagdad [Internet]*. 2022 Oct. 17 [cited 2022 Nov. 15];64(3):195-8. Available from: <https://iqjmc.uobaghdad.edu.iq/index.php/19JFacMedBaghdad36/article/view/1957>

تقدير دور Malondialdehyde في بدء وتطور اعتلال الشبكية السكري

د. زينة محمد حسن / كلية الطب / قسم الكيمياء / جامعة بغداد
أ.م.د. رنا علي حمدي كلية الطب / قسم الكيمياء / جامعة بغداد
د. ابتهاج نوري البصام / مركز السكري التخصصي للغدد الصم

الخلاصة

الخلفية: النتيجة الأكثر شيوعاً لمرض السكري هي اعتلال الشبكية السكري. يلعب الإجهاد التأكسدي جزءاً مهماً من مسببات مرض السكري واعتلال الشبكية السكري. مالونديالدهيد هو دايالدهيد ذو سمية معتدلة وكثيراً ما يستخدم كمؤشر حيوي للإجهاد التأكسدي وبيروكسيد الدهون.

الأهداف: كان الهدف من هذه الدراسة هو تقييم مستويات مصطلح الدم للمالونديالدهيد في المرضى الذين يعانون من اعتلال الشبكية السكري والذين يعانون من مرض السكري النوع الثاني. ومقارنةً مستواه بالأصحاء. بالإضافة إلى ذلك، التحقق من العلاقة بين مستوى مالونديالدهيد و فحص السكر التراكمي في الدم.

الموضوعات والطرق: شملت دراسة الحالات والشواهد هذه (120) فرداً تتراوح أعمارهم بين 40 و 70 عاماً. تم إنشاء ما مجموعه 3 مجموعات من الأشخاص: 40 من مرضى السكري من النوع 2 المصابين باعتلال الشبكية كانوا جزءاً من المجموعة 1، 40 كان مرضى السكري من النوع 2 غير المصابين باعتلال الشبكية جزءاً من المجموعة 2، وكان 40 شخصاً من المجموعة الضابطة.

النتائج: كانت مستويات مصطلح الدم المالونديالدهيد أعلى بشكل ملحوظ في مرضى السكري الذين يعانون من اعتلال الشبكية وبدونه مقارنةً بالضوابط ($p = 0.0001$)، وكذلك في مرضى السكري الذين يعانون من اعتلال الشبكية عند مقارنتهم بمرضى السكري غير المصابين باعتلال الشبكية ($p = 0.0001$). بالإضافة إلى ذلك، في مجموعة مرضى السكري الذين يعانون من اعتلال الشبكية، تم اكتشاف علاقة إيجابية ذات دلالة إحصائية بين مستوى مصطلح دم المالونديالدهيد والدم و HbA1C ($r=0.931$ ، $p=0.0001$).

الخلاصة: ارتفاع مستويات مصطلح الدم المالونديالدهيد هو مؤشر على زيادة بيروكسيد الدهون التي قد تكون متورطة في مسببات أمراض اعتلال الشبكية في مرضى السكري من النوع 2 غير المنضبط.

الكلمات المفتاحية: الإجهاد التأكسدي، اعتلال الشبكية السكري، المالونديالدهيد