

The Advantages of Combined Regional and General Anesthesia in Ophthalmic Surgery in Children

Doi: <https://doi.org/10.32007/jfacmedbagdad.2063>

Rand S. Abdul-Sattar Al-Ani* FICMS (Ans. & ICU)
Ali H. Mosleh Al-Maini* FICMS (Ans. & ICU)



This work is licensed under a [Creative Commons Attribution-NonCommercial 4.0 International License](https://creativecommons.org/licenses/by-nc/4.0/).

Abstract

Background: Regional anesthesia is a common procedure in an adult patient undergoing ophthalmic surgery, but it cannot be done alone in the pediatric age group. General anesthesia is accompanied by complications intra- and post-operatively.

Objectives: to determine whether or not using regional anesthesia in combination with general anesthesia in pediatric eye procedures improves patient outcomes.

Methods: Forty children, with an age range of 6 - 12 years were included in the study that was conducted at Ghazi Al-Hariri Hospital / Department of Ophthalmology and Ibn Al-Haytham (Ophthalmology Hospital), both teaching hospitals, from December 2018 to October 2019. These children were allocated into one of two groups: GA (general anesthesia) and GA-R (general anesthesia-regional anesthesia). Heart rates, mean arterial blood pressure, oculo-cardiac reaction, and postoperative nausea and vomiting were measured. All required approvals were obtained from the scientific committee of the Iraqi Board for Medical Specializations. Statistical analysis was done using SPSS V26, with a P value of <0.05 considered significant.

Results: None of the patients developed oculo-cardiac reactions or needed additives to the anesthesia given during surgery in the GA-R compared to the GA group. Intra-operative measurements of heart rates and mean arterial blood pressure were lower in the GA group than in the GA-R group ($p < 0.05$). Compared to the GA group, the GA-R group had a lower incidence of postoperative nausea and vomiting ($p < 0.05$). More patients in the GA group needed analgesia than in the GA-R group.

Conclusion: Using regional anesthesia as a peribulbar block with general anesthesia is a safe and successful procedure in pediatric ocular surgeries.

Keywords: Regional Anesthesia, General Anesthesia, Peribulbar block, Oculo-cardiac Reaction

J Fac Med Baghdad
2023; Vol.65, No. 3
Received: Feb. 2023
Accepted: Sept. 2023
Published: Oct. 2023

Introduction:

Ophthalmic surgery is a frequent treatment performed on children, and the conventional method of pain management is general anesthesia (1). Consequences, both intra- and post-operative, associated with various anesthetics complication, including the oculo-cardiac reflex (OCR), postoperative nausea and vomiting (PONV), postoperative irritation, and discomfort, are serious issues in juvenile ocular procedures. Additionally, repeated sobbing and squinting might be harmful to the intraocular pressure (IOP) and the corneal sutures (2, 3). The OCR is a trigeminal vagal response that is triggered by tension on the extraocular muscles, compression mostly on the orbit, trauma, or an orbital hematoma (4). Other triggers of the OCR include hypercarbia and cerebral hypoxia. Mild anesthesia may also play a role (6). During strabismus surgery, there has been a documented incidence of transient cardiac arrest of around 1 in 2200 patients (7).

There have been reports of arrests in the sinoatrial node and fibrillation in the ventricular node (8, 9). PONV is a common problem following ophthalmic surgery. Repeated vomiting can affect the integrity of corneal sutures, increase the cost because of the use of antiemetic drugs, and prolong the postoperative stay. PONV is mediated by the oculo-emetic pathway, which shares the afferent limb of the reflex arc with OCR (11-12).

A double injection peribulbar block helps lower postoperative pain in patients who have had ocular surgeries (3, 13). Since peribulbar block generates a hypotonic eye, it is not often employed in strabismus surgery (14).

However, regional anesthesia is a common procedure in adult patients undergoing ophthalmic surgery, but it cannot be done alone in the pediatric age group. Double injection peribulbar block as an adjuvant to general anesthesia has been investigated and reported to be effective. However, this technique is risky as it is administered invisibly and may result in globe perforation and intravascular injection.

The double injection peribulbar block in children carries potentially more risks than in adults because there are important anatomical differences between

* Corresponding Author: Ghazi Al-Hariri Hospital.

dr.randsadi@yahoo.com

**Al-Mustansiriya University, College of
Medicine.alimaini2000@gmail.com

the two age groups. The eye size relative to the bony orbit is much greater in childhood. The eye occupies almost 50% of the volume of the bony orbit at birth and 33% at four years, while the adult eye takes up only 22% of the orbital volume.(1-15-18) There is little evidence concerning its usage in the pediatric age group. Hence, the recent study attempted to overcome these complications by using regional anesthesia as an adjuvant to general anesthesia in ophthalmic surgery in children.

Patients and Methods:

A randomized clinical, non-blinded study was conducted at Ghazi Al-Hariri Hospital / Department of Ophthalmology and Ibn Al-Haytham (Ophthalmology Hospital), both teaching hospitals, from December 2018 to October 2019. Forty children with an American Society of Anesthesiology (ASA) Classification I or II (19) undergoing elective ophthalmic surgeries were enrolled. The institutional ethical committee's approval of the Scientific Council of Anesthesia and Intensive Care of the Iraqi Board was obtained for the study. Parents' consent was given to all patients, including their agreement on the type of anesthesia used.

Two types of ophthalmic surgeries were performed: Intraocular (cataract) or extraocular (strabismus).

Inclusion criteria: Children aged 6-12 years, ASA I or II, elective ophthalmic surgeries, and body weight between 15-40 Kg.

Exclusion criteria: Family refusal, prolonged surgery >2 hours, endophthalmitis, eye injury, cerebral palsy, psychological problems, single eye, congenital abnormalities, heart diseases, and drug allergy.

Patients were randomly divided into two groups: Those who received general anesthesia only (GA group, n=20) and those who received combined general and regional anesthesia (single injection peribulbar block anesthesia) (GA-R group, n=20).

Post-operative pain was recorded using a VAS score. (20)

Post-operative recovery was assessed using a modified Aldrete Score in pediatrics (21) every 5 minutes, and the time to achieve the full score of 9 was recorded. The occurrence of PONV was assessed for 12 hours postoperatively. The complications due to peribulbar block were assessed. The data was analyzed using SPSS version 25. The data was presented as mean, standard deviation, or frequencies, as appropriate. The independent t-test (two-tailed) was used to compare the means of continuous variables between the study groups. A P value < 0.05 was considered significant.

Results:

There were no significant differences in the demographic characteristics between the two groups (table 1).

Table 1: Demographic and operative variables of the two study groups

Variable	GA group no.=20	GA-R group no.=20
Age (years) - Mean ± SD	7.9±1.4	8.3±1.8
Weight (Kg) - Mean ± SD	25.6±6.5	28.7±5.2
Sex (male/female) - (No.)	10/10	8/12
Type of surgery - (No.)		
Strabismus	11	12
Cataract	9	8
Duration of surgery (minutes) - Mean ± SD	62.1±12.2	59.4±11.6

Among the GA-R group, not a single patient exhibited OCR, in contrast to 12 children among the GA group (60%) (p<0.05). There was a need for an atropine prescription for five of them.

None of the GA-R group required intraoperative supplemental fentanyl. In contrast, five patients (25%) in the GA group needed only one dose of fentanyl (p< 0.05).

The mean end-tidal sevoflurane concentrations of the two groups were very close (2.4 ±0.39 vs 2.6± 0.41). At T0 and T1, both groups' Mean arterial pressure (MAP) and heart rate (HR) values were not statistically significant. In comparison to both its base level and the GA-R group, the values at (T2–T6) for the GA group were statistically substantially higher (p< 0.05). (Figures 1 and 2) and (Tables 2 and 3)




Figure 1: Intra-operative MAP in the study groups

Table 2: Mean intraoperative arterial pressure by time and group

Time	GA group no.=20 Mean ± SD	GA -R group no.=20 Mean ± SD	P-value
T0	90.0 ± 6.43	88.7 ± 7.45	0.350
T1	84.2 ± 4.66	86.6 ± 7.14	0.053
T2	87.0 ± 0.23	87.1 ± 0.16	0.028
T3	86.9 ± 0.20	87.1 ± 0.14	0.007
T4	86.9 ± 0.21	87.1 ± 0.16	0.008
T5	86.9 ± 0.24	87.1 ± 0.19	0.024
T6	86.8 ± 0.23	87.1 ± 0.16	0.006




Figure 2: Intra-operative HR in the study groups

Table 3: Mean intraoperative heart rate by time and group

Time	GA group n.=20 Mean ± SD	GA -R group n.=20 Mean ± SD	P-value
T0	89.6 ± 7.3	91.3 ± 14.37	0.562
T1	95.1 ± 6.29	97.2 ± 15.05	0.541
T2	97.1 ± 0.18	96.8 ± 0.22	0.016
T3	97.0 ± 0.13	96.8 ± 0.20	0.015
T4	97.1 ± 0.19	96.9 ± 0.21	0.023
T5	97.9 ± 0.18	96.9 ± 0.26	0.016
T6	97.1 ± 0.16	96.8 ± 0.23	0.006

Table 4: Postoperative comparison of the two study groups

Variable	GA group no.=20	GA-R group no.=20	p-value
Time to achieve full Aldrete score (minutes)	76 ± 0.44	24 ± 0.62	0.001
Time to first analgesic (hour)	0.36 ± 0.56	1.14 ± 0.72	0.001
Number of cases needing analgesia within the first 12 hours	16 (80%)	8 (40%)	0.0099
Number of analgesic doses administered within the first 12 hours	1 14 (70%)	6 (30%)	0.0114
	2 10 (50%)	4 (20%)	0.0466
Frequency of PONV during the first 12 hours	4 (20%)	0	0.0348
Total number of children needing antiemetic	8 (40%)	2 (10%)	0.0285

Discussion:

The current research demonstrated that a peribulbar block administered via a single dose injection decreased the risk of OCR, patients undergoing fentanyl necessity, and PONV because once taken in conjunction with general anesthesia in pediatric ophthalmic procedures. Additionally, it lengthened the time until the patient required their first postoperative analgesic medication.

Earlier research (1, 22) in pediatrics employed instruments that were 16 millimeters long while doing double injections of peribulbar blocks. The injection method used in this research was carried out using a relatively short needle, measuring 12 millimeters in length and having a gauge size 28. This was done to decrease the risks of side effects associated with longer needles, including bleeding or lacerations of the cornea (23). Using a single injection in the current research wasn't associated with the health problems encountered during the double injection system. These complications noted in the double injection method are severe but uncommon, such as Injuries to the vascular system or instances of misdirection (23).

In the post-anesthesia care unit (PACU) setting, the time it took for either group to get a complete Aldrete score was not similar (76 ± 0.44 minutes in the GA-R group and 24 ± 0.62 minutes in the GA group).

The incidence of PONV during 12 hours post-operative and the number of children who required anti-emetics were significantly lower in the GA-R group [4 (20%) and 0 (0%), respectively, than the GA group 8 (40%) and 2 (10%), respectively], ($P < 0.05$).

Throughout the first 12 postoperative hours, significantly fewer children in the GA-R group needed analgesics than those in the GA group ($P < 0.05$). When compared to the GA group, the GA-R group had a much longer period before needing their first analgesia ($p = 0.001$), and they also required less analgesia treatment ($P < 0.05$), Table 4.

Surgeons prefer the peribulbar block because the eyeball becomes centrally located after the block, which is beneficial in strabismus surgery and is associated with a high risk of OCR.

Compared to the GA category, the frequency of OCR was much lower in the GA-R due to the peribulbar block. This may be related to limiting the sensory limb of OCR and limiting the sensory inputs from muscular tension, which is similar to the results of published research (24, 25).

In accordance with our results, Deb et al. (1) reported that 4% of children in the peribulbar block group experienced OCR compared to 60% in the GA group when they studied the efficacy of a double injection technique for peribulbar block as adjuvant to GA in ophthalmic surgery.

The research carried out by Gupta et al. (22) found that the use of double injection peribulbar block in combination with GA reduced OCR occurrence in people undergoing strabismus surgical procedures. The percentage of children who experienced OCR was 13% in the GA-R group and 94% in the GA group.

The increased occurrence of oculocardiac reflex in the GA category found in the latter research compared to the current investigation may be attributable to the younger age group (2 – 13 years) since young children have a significant propensity to have higher resting vagal tone than adults (26). In addition, all participants were about to receive a strabismus operation, a procedure recognized to result in a greater rate of OCR (27, 28).

In the present study, the peribulbar block administered with a single injection was sufficient to provide appropriate intraoperative analgesia, as shown by decreased intraoperative fentanyl required and maintained hemodynamics in the GA-R group. These findings are consistent with those of earlier studies that looked into localized blocks' intraoperative analgesic impact during ocular surgical procedures (1, 3, and 29). Using a peribulbar block with a single injection as just an adjuvant to GA during the present trial helped reduce the likelihood of PONV. This was in agreement with the results of Gupta et al. (22), who found that using a double injection peribulbar block satisfactorily significantly reduced the PONV in people who had undergone ophthalmic surgical operation while having received GA. The participants in these studies were given GA while having the operation.

Adolescent patients with vitreoretinal procedures who were given a peribulbar block by a single injection in combination with GA had outcomes comparable to those described above (30). Research on a single-injection peribulbar block coupled with general anesthesia in children having ocular operations was published by Elgohary et al. and reached the same findings (31).

Subramaniam et al. (3) reported the rate of PONV to be 54.8% in patients who received double injection peribulbar block, compared to 81.4% in those who had GA. They explained this high incidence across both groups by the longer duration of the vitreoretinal operation and the diminishing effect of the block.

While looking at the peribulbar block both as an adjuvant to GA and GA by itself in ocular procedures, Chhabra et al. (14) and Moral et al. (32) were unable to find a difference in PONV, which is inconsistent with our results. The utilization of total intravenous anesthesia (TIVA) may have contributed to this result owing to the antiemetic effects of propofol (33).

Although Shende et al. (24) evaluated the use of the peribulbar block as an adjuvant to GA in surgeries for retinal detachment, researchers suggested that perhaps the duration of the first rescue analgesia was equal in the peribulbar and GA groups.

This finding is by results reported by Ghali and Btarny in 2010 (32) when they studied the effect of single injection peribulbar block in adults undergoing vitreoretinal. Also, Subramaniam et al. (3) reported similar results when using double injection peribulbar block in pediatrics.

In contrast, when Shende et al. (24) investigated the usage of the peribulbar block as an adjuvant to general anesthesia during retinal detachment surgery, they found that the time first to rescue analgesia was comparable in peribulbar and GA groups. This could be explained by using a lower concentration of bupivacaine (0.25%) in the peribulbar block group and morphine in the GA group, as stated by Ghali and Btarny (32).

Conclusion:

Utilizing a local single-injection peribulbar anesthetic drug with GA in pediatric ocular surgeries is a useful method that may be used as an alternate to GA alone. It results in a low frequency of opioid-related complications, fewer requests for intraoperative opioids, steady intraoperative hemodynamics, fewer instances of nausea and vomiting after surgery, and enhanced postoperative analgesia.

Authors' contributions:

Dr. Rand Saadi Abdul-Sattar Alani: Writing the project, collecting data, writing draft, and research.

Dr. Ali Hadi Mosleh Al-Maini: Supervisor, concept of the study, reviewing manuscript

Authors' Declaration

Conflicts of Interest: None.

We hereby confirm that all the Figures and Tables in the manuscript are ours. Besides, the Figures and images, which are not ours, have been permitted re-publication and attached to the manuscript. Authors sign on ethical consideration's approval-Ethical Clearance: The project was approved by the local ethical committee of (the Iraqi Board for Medical Specializations) according to code number (No 88624.2.2020).

References

1. Deb K, Subramaniam R, Dehran M, Tandon R, Shende D. Safety and efficacy of peribulbar block as adjuvant to general anaesthesia for paediatric ophthalmic surgery. *Paediatric Anaesthesia* 2001; 11:161–7. DOI: 10.1046/j.1460-9592.2001.00623.x
2. Kumar C, Dodds C, Gayer S. *Oxford Specialist Handbooks in Anaesthesia Ophthalmic Anaesthesia* 2012, p. 189–224. https://www.researchgate.net/publication/31164882_8_Oxford_Handbook_of_Ophthalmic_Anaesthesia
3. Subramaniam R, Subbarayudu S, Rewari V, Singh R, Madan R. Usefulness of pre-emptive peribulbar block in pediatric vitreoretinal surgery: a prospective study. *Regional Anesthesia and Pain Medicine* 2003; 28:437. DOI: 10.1053/rapm.2003.50032
4. Breijo Marquez FR. *Abnormal Heart Rhythms*. 2015;97:105. DOI: 10.5772/59212
5. Hahnenkamp K, Honemann CW, Fischer LG, Durieux ME, Muehlendyck H, Braun et al. Effect of different anaesthetic regimes on the oculo-cardiac

- reflex during paediatric strabismus surgery. *Paediatr Anaesth.* 2000; 10:601–8. DOI: 10.1111/j.1460-9592.2000.00588.x
6. Gilani MT, Sharifi M, Najafi MN, Mashhadi MGE. Oculo-cardiac reflex during strabismus surgery. *Reviews in. Clinical Medicine.* 2016; 3:4–7. <https://doi.org/10.17463/RCM.2016.01.002>
7. Bietti OB. Problems of anesthesia in strabismus surgery. *Int Ophthalmol Clin.* 1966; 6:727–37. <https://journals.lww.com/internatophthalmology/Citation/1966/00630/>
8. Chowdhury T, Schaller B, Koerbe A. Functional Consequences of the Trigemino-cardiac Reflex 2015; 14:169:213. DOI: 10.1097/ANA.0000000000000065
9. Smith RB, Douglas H, Petruscak J. The oculo-cardiac reflex and sino-atrial arrest. *Can Anaesth Soc J.* 2002; 19:138–42. https://www.jidmr.com/journal/wpcontent/uploads/2018/04/47D17_453_Srikanth_Gadicherla.pdf
10. Sadhasivam S, Shende D, Madan R. Prophylactic ondansetron in prevention of postoperative nausea and vomiting following pediatric strabismus surgery: a dose response study. *Anesthesiology* 2000; 92:1035–42. DOI: 10.1097/00000542-200004000-00021
11. Watcha MF. The cost/effective management of postoperative nausea and vomiting. *Anesthesiology* 2000; 92:931–3. DOI: 10.1097/00000542-200004000-00007
12. Wolf R, Morinello E, Kestler G, Käsmann-Kellner B, Bischoff M, Hager T, et al. PONV after strabismus surgery: Risk adapted prophylaxis, 2016 Jul;65(7):507-13. DOI: 10.1007/s00101-016-0183-2
13. Kristin N, Schonfeld CL, Bechmann M, Bengisu M, Ludwig k, Scheider A, et al. Vitreoretinal surgery: pre-emptive analgesia. *Br J Ophthalmol* 2001; 85:1328–31 DOI: 10.1136/bjo.85.11.1328
14. Chhabra A, Pandey R, Khandelwal M, Subramaniam R, Gupta S. Anesthetic techniques and postoperative emesis in pediatric strabismus surgery. *Reg Anesthesia Pain Med* 2005; 30(1):43–7. DOI: 10.1016/j.rapm.2004.08.023
15. Coté CJ, Lerman J, and Anderson BJ. *A Practice of Anesthesia for Infants and Children* 2018; 34:790:810. <https://www.worldcat.org/title/A-practice-of-anesthesia-for-infants-and-children/oclc/225874385>
16. Ghali B, Ram J, Makkar JK, Wig J, Kaushik S. Subtenon block compared to intravenous fentanyl for perioperative analgesia in pediatric cataract surgery 2009; 108 (4):1132–8. DOI: 10.1213/ane.0b013e318198a3fd
17. Parulekar MV, Berg S, Elston JS. Adjunctive peribulbar anesthesia for paediatric ophthalmic surgery: are the risks justified? *Paediatric Anaesth* 2002; 12:85–6. DOI: 10.1046/j.1460-9592.2002.0787a.x
18. Forrester JV, Dick AD, McMenamin PG, Roberts F, Pearlman E. *The Eye: Basic Sciences in Practice Fourth Edition* 2016; 1:1:60. <https://doi.org/10.1016/C2012-0-07682-X>
19. Jennings D, Chernega J. *Emergency Guide for Dental Auxiliaries* 2012; 2:23-233. <https://doi.org/10.1016/C2012-0-07682-X>
20. Geyer H, Kaufman DM, Milstein MJ, Kaufman's *Clinical Neurology for Psychiatrists* 2016; 315:688. <https://www.asia.elsevierhealth.com/kaufmans-clinical-neurology-for-psychiatrists-e-book-9780323461313.html>
21. Bissonnette B, Lockman JL and Easley RB. *Pediatric Anesthesia Basic Principles-State of the Art-Future, Emergence and Postoperative Care* 2011; 58:924:939 https://books.google.iq/books/about/Pediatric_Anesthesia.html?id=RYe4GBD6LrOC&redir_esc=y
22. Gupta N, Kumar R, Kumar S, Sehgal R, Sharma KR. Prospective randomized double-blind study to evaluate the effect of peribulbar block or topical application of local anaesthesia combined with general anaesthesia on intra-operative and postoperative complication during pediatric strabismus surgery. *Anaesthesia* 2007; 62:1110–3. DOI: 10.1111/j.1365-2044.2007.05220.x
23. Ripart J, Lefrant JY, L'Hermite JI, Borzli F, Nouvellon E, Fabbro-Peray P, et al. Caruncle single injection episcleral (sub-tenon) anesthesia for cataract surgery: mepivacaine versus a lidocaine-bupivacaine mixture. *Anesth Analg* 2000; 91:107–9. DOI: 10.1097/00000539-200007000-00020
24. Shende D, Sadhasivam S, Madan R. Effect of peribulbar bupivacaine as an adjuvant to general anesthesia on postoperative outcome following retinal detachment surgery. *Anesthesia* 2000; 55:970–5. DOI: 10.1046/j.1365-2044.2000.01535.x
25. Dehghani A, Montazeri K, Masjedi A, Karbasi N, Ashrafi L, Saeedian et al. Adjunctive Local Application of Lidocaine during Scleral Buckling under General Anesthesia 2011; 177:181. <https://pubmed.ncbi.nlm.nih.gov/22454732/>
26. Shalini G, Chakrapani K, Harsha N, Karanam S, Nambula S. Occurrence of Oculo-cardiac Reflex During Strabismus Surgery Vol. 30, No. 1, July-September 2019; 28:31. https://journals.lww.com/djo/Abstract/2019/30010/Occurrence_of_Oculo_Cardiac_Reflex_During.6.aspx
27. Gilani MT, Sharifi M, Najafi MN, Mashhadi MGE. Oculo-cardiac reflex during strabismus surgery. *Reviews in. Clinical Medicine.* 2016; 3:4–7. <https://doi.org/10.17463/RCM.2016.01.002>
28. Ha SG, Huh J, Lee BR, Kim SH. Surgical factors affecting oculo-cardiac reflex during strabismus surgery, 2018; 18:103:1-5. <https://doi.org/10.1186/s12886-018-0771-9>
29. Chhabra A, Sinha R, Subramaniam R, Chandra P, Narang D, Garg SP, et al. Comparison of sub-Tenon's block with i.v. fentanyl for paediatric vitreoretinal surgery. *Br J Anesthesia* 2009; 103(5):739–43. DOI: 10.1093/bja/aep230
30. Ghali AM, Hafez A. Single-injection percutaneous peribulbar anaesthesia with a short needle as an alternative to the double injection

technique for cataract extraction. *Anesth Analg* 2010; 110(1):245-7.

DOI: 10.1213/ANE.0b013e3181c293bd

31. Elgohary MM, Sherif H. Single-injection peribulbar block combined with general anesthesia in children undergoing ophthalmic surgery. *Egyptain J Anaesthesia* 2011; 77-82.

<https://doi.org/10.1016/j.egja.2011.02.003>

32. Morel J, Pascal J, Charier D, De Pasquale V, Gain P, Auboyer C, et al. Preoperative peribulbar block in patients undergoing retinal detachment surgery under general anesthesia: a randomized double-blind study. *Anesthesia Analgesia* 2006; 102:1082-7.

DOI:10.1213/01.ane.0000198638.93784.42

33. Moghaddam AAS. Assessment of the Rate of Nausea, Vomiting, and Pain in Patients with Strabismus Anaesthetised with Propofol. Volume 7, Number 2, 2005; 30:32.

https://www.researchgate.net/publication/242216724_Assessment_of_the_Rate_of_Nausea_Vomiting_and_Pain_in_Patients_with_Strabismus_Anaesthetised_with_Propofol

34. Ghali AM, El Btarny AM. The effect on outcome of peribulbar anaesthesia in conjunction with general anesthesia for vitreoretinal surgery. *Anaesthesia* 2010; 65:249-53.

DOI: 10.1111/j.1365-2044.2009.06191.x

35. Salman IA. Effects of halothane and isoflurane on uterine muscle relaxation during caesarian section. *JFacMedBagdad [Internet]*. 2013 Jan. 2 [cited 2023 Jan. 23]; 54(4):314-6.

<https://doi.org/10.32007/jfacmedbagdad.544692>

36. Salman IA, Kanaan SA. Comparison between the effect of propofol (TIVA) and inhalational anesthesia (halothane) on early postoperative nausea and vomiting. *JFacMedBagdad [Internet]*. 2013 Jul. 1 [cited 2023 Jan. 23]; 55(2):111-4. DOI:

<https://doi.org/10.32007/jfacmedbagdad.552637>

How to cite this Article

Abdul-Sattar RS, Al-Maini AH. The Advantages of Combined Regional and General Anesthesia in Ophthalmic Surgery in Children. *JFacMedBagdad [Internet]*. [cited 2023 Sep. 28];65(3):173-8. Available from: <https://iqjmc.uobaghdad.edu.iq/index.php/19JFacMedBaghdad36/article/view/2063>

Copyright (c) 2023 Rand Saadi Abdul-Sattar, Ali Hadi Al-Maini, Mais Razaq Jaafar

مزايا الجمع بين التخدير الموضعي والعام في جراحة العيون عند الاطفال

د. رند سعدي عبدالستار العاني: طبيبة اختصاص تخدير وعناية مركزة، مستشفى غازي الحريري
الأستاذ المساعد علي هادي المعيني: استاذ في التخدير والعناية المركزة، الجامعة المستنصرية/ كلية الطب
الخلاصة

الخلفية: التخدير الموضعي هو إجراء شائع لدى المرضى البالغين الذين يخضعون لجراحة العيون ولكن لا يمكن القيام به بمفرده في الفئة العمرية للأطفال. ويرافق التخدير العام مضاعفات أثناء الجراحة وبعد العملية الجراحية.

الهدف: تحديد ما إذا كان استخدام التخدير الموضعي مع التخدير العام في إجراءات عيون الأطفال يحسن نتائج المرضى أم لا.
الطريقة: تم شمول أربعين طفلاً تتراوح أعمارهم بين ستة أعوام وأثني عشر عامًا في الدراسة التي أجريت في مستشفى غازي الحريري/ قسم العيون ومستشفى ابن الهيثم (مستشفى العيون) وكلاهما مستشفى تعليمي، في الفترة من كانون الأول 2018 إلى تشرين الأول 2019، وقسموا إلى مجموعتين: GA (التخدير العام)، و GA-R (التخدير العام - التخدير الموضعي). تم قياس معدلات ضربات القلب (HR)، متوسط الضغط الشرياني (MAP)، تفاعل القلب (OCR) والغثيان والقيء بعد الجراحة (PONV). تم الحصول على جميع الموافقات المطلوبة من المجلس العراقي للتخصصات الطبية. تم إجراء التحليل الإحصائي باستخدام SPSS V26. تعتبر قيمة $P < 0.05$ ذات أهمية إحصائية.

النتائج: لم يصب أي مريض بتفاعل عيني قلبي أو إضافات ضرورية للتخدير أثناء الجراحة في GA-R مقارنة بمجموعة GA. كانت القياسات أثناء العملية لدقات القلب ومتوسط الضغط الشرياني قراءة أقل في مجموعة GA، مجموعة GA-R ($p < 0.05$). بالمقارنة مع مجموعة GA، فإن مجموعة GA-R قد قللت من حدوث الغثيان والقيء بعد الجراحة $p < 0.05$. واحتاج عدد أكبر من المرضى في مجموعة GA إلى تسكين أكثر من مجموعة GA-R.

الاستنتاج: يعد استخدام التخدير الموضعي كحقن حول المقلة مع التخدير العام إجراءً آمنًا وناجحًا في جراحات عيون الأطفال.
الكلمات المفتاحية: التخدير الموضعي، التخدير العام، تخدير محيط العين، منعكس عيني قلبي.