

Platelet Parameters in Nonthrombocytopenic Preeclampsia: A Case-Control Study

Doi: <https://doi.org/10.32007/jfacmedbagdad.2084>

Ahmed J. Alhusaynei* DOG/ FICOG
Hanan Dh. Nayyef* MBChB



This work is licensed under a [Creative Commons Attribution-NonCommercial 4.0 International License](https://creativecommons.org/licenses/by-nc/4.0/)

Abstract

Background: Preeclampsia is a pregnancy-specific, multisystem condition characterized by the onset of de novo hypertension and proteinuria occurring in a previously normotensive women after the twentieth week of pregnancy. Pregnancy is associated with physiological adaptation led to change in hematological system including platelet parameters.

Objectives: Is to compare platelet count, platelet indices, namely mean platelet volume platelet distribution width and platelet count to mean platelet volume MPV ratio in preeclamptic patients with normal pregnant women.

Patients & methods: A case-control study was carried out in obstetrics and gynecology department at Al-Khansaa maternity and childhood Teaching hospital, Mosul, Iraq over a period extending from February 2020 to the end of December 2020. A sample of 120 pregnant women participated in the study fulfilling inclusion criteria. Sixty of them diagnosed with preeclampsia PE (30 pregnant women with mild PE and 30 pregnant women with severe PE) and sixty healthy pregnant women as control group. Platelet count, mean platelet volume and platelet distribution width (PDW) were measured and statistical analysis by mini tab program was done for all collected data.

Results: A significant linear correlation between mean platelet volume (MPV) and platelet distribution width (PDW) with mean arterial pressure in Preeclampsia group with p value of (0.025 and 0.001) respectively and a significant inverse correlation between the platelet count and mean arterial pressure with p value of (<0.001) was observed.

Conclusion: Platelet parameters are cost-effective, easily measurable, and practical markers which can be used for monitoring of preeclampsia. Mean platelet volume (MPV) and platelet distribution width (PDW) has a significant linear correlation with mean arterial pressure in severe preeclampsia and can be used as biomarker for severity of preeclampsia in high-risk pregnant women.

Keywords: mean platelet volume; platelet count; platelet distribution width.; platelet parameters; preeclampsia thrombocytopenia.

J Fac Med Baghdad
2023; Vol.65, No.4
Received: Mar., 2022
Accepted: July, 2023
Published: Jan.2024

Introduction:

Preeclampsia is an idiopathic disorder of pregnancy characterized by proteinuria and hypertension (1), the disorder is specific to pregnancy characterized by new onset of hypertension and proteinuria after 20 weeks of gestation. Preeclampsia usually presents during pregnancy; however, it may sometimes manifest in the postpartum period in a previously normotensive woman. Preeclampsia is defined as blood pressure of greater than 140 mmHg systolic and/or 90

mmHg diastolic, measured on at least two occasions, not less than four hours apart in the presence of at least 300mg protein in a 24-hour collection of urine or urinary protein / creatinine ratio of greater than 30mg/mmol (2,3). Hypertension complicates 6–12% of all pregnancies and includes two relatively benign conditions (chronic and gestational hypertension) and the more severe conditions of preeclampsia or eclampsia (4). Preeclampsia complicates 5% of all pregnancies worldwide (5). Platelets count (PC) and mean platelet volume (MPV) are largely unchanged in most pregnant women, even though platelets survival is reduced. Platelet reactivity is increased in the second and third trimesters and

*Corresponding Author: Al-Khansaa Maternity Teaching Hospital/ Al-Mosul

* ajm@uomosul.edu.iq
thanonhanan@gmail.com

does not return to normal until about 12 weeks after the delivery (4). Platelets play an important role in the pathogenesis of preeclampsia. The platelets are normally present in the blood stream in an inactive state, but they can be activated instantly when they come in contact with the damaged or activated endothelial wall (6). Increased consumption of platelets as well as decreased life span in the uteroplacental circulation has been suggested to be the explanation of the reduction in the number of circulating platelets during pregnancy. Platelet counts as well as other hemostatic factors return to normal after 6 weeks of delivery (7). The platelets count decreases by an average of 10% during the third trimester of uncomplicated pregnancy due to hemodilution (8,9). Platelets count is always included in the reporting of complete blood count results. Automated blood cell analyzers have measured mean platelet volume (MPV) and platelet distribution width, and as a result the PC to MPV ratio can be calculated. The MPV is a measure of the average volume of platelets in a sample. Large platelets are often biochemically more active than smaller platelets; high mean platelet volume is associated with less bleeding in patients with severe thrombocytopenia (10). In healthy people, there is an inverse relationship between platelet count and size. Because of this inverse relationship, the MPV and the platelet count must be considered together. The increased MPV and increased platelet aggregation have been defined as precursors to the onset of preeclampsia and intrauterine growth restriction (11,12). The MPV is elevated proportionally with the severity of preeclampsia (13). When platelets are activated, there will be an increase in production and they become larger in size which causes increased platelet indices such as mean platelet volume (MPV). So, platelet indices can give an idea of platelet activation (14). Platelet distribution width (PDW) is a measurement of platelet anisocytosis (12,15). PDW can act as an indicator of platelet volume variability and increases in the presence of platelet anisocytosis. In addition, PDW can be affected by the morphology change of platelet when the platelet is activated during inflammation and thrombosis process (16). Platelet distribution width is considered a marker of platelet function and activation (17). On the other hand, lowered platelet count and increased platelet indices in pregnant women during antenatal check may be considered as a risk factor for development of preeclampsia and eclampsia. This group of pregnant women may be considered for special attention to prevent the development of preeclampsia (18,19).

Patients and method

A case-control study was carried out in obstetrics and gynecology department at Al-Khansaa maternity and childhood Teaching hospital, Mosul, Iraq over a period extending from February 2020 to the end of December 2020.

Inclusion criteria for cases and controls:

- Viable singleton pregnancy.
- Third trimester of pregnancy.

Exclusion criteria for cases and controls:

- Chronic hypertension.
- Anemia
- Cardiovascular disease.
- Diabetes mellitus type 1 and 2.
- Multiple gestations.
- Smoking.
- Chronic renal disease.
- Intrauterine fetal death.
- Fetal anomalies.
- Autoimmune disease.
- Eclampsia, HELLP syndrome (hemolysis, elevated liver enzyme and low platelet).
- Thrombocytopenia (platelets count less than $150 \times 10^9/L$).
- Maternal age not less than 16 years and not more than 35 years.

A sample of 120 pregnant women participated in the study fulfilling inclusion criteria, who presented at antenatal care out-patient clinic and/or admitted in maternity wards. Sixty of them diagnosed with preeclampsia PE (30 pregnant women with mild PE and 30 pregnant women with severe PE) and sixty healthy pregnant women as control group. Cases with severe PE were using aspirin from early pregnancy and stopped uses at 36 weeks of pregnancy. The diagnosis of preeclampsia was made by NICE guideline criteria (20); Include clinical features, blood pressure measurement and laboratory investigation which included a complete blood count, renal function test, liver function test and detection of protein in urine done by clean-catch midstream or catheter specimens of collected urine > 4 hours apart with 2+ on reagent strip, urine dipstick testing 1+ = 0.3 g/L, 2+ = 1 g/L and 3+ = 3 g/L.

Platelet count, mean platelet volume and platelet distribution width were measured and statistical analysis was done for all the data collected.

Blood pressure was measured by auscultatory method with a properly calibrated standard mercury sphygmomanometer, the woman should be rested and sitting at a 45-degree angle or in left lateral position with the cuff on the left arm at the level of the heart. Pulse pressure (PP) was calculated as the difference between SBP and DBP. Mean arterial pressure (MAP) was calculated as DBP plus 1/3 of PP. hematological analyzer Abbott Diagnostic Division (Max - Plank -Ring 2) used

for determination of the hemoglobin, platelet count (PC), platelet indices, which includes mean platelet volume and platelet distribution width as a part of full blood count. MPV was recorded as femtoliter (fL). The range of normal values for platelet count in our laboratory was (150-400×10⁹ /L), for mean platelet volume 7.6-10.6 fL and for platelet distribution width 14-19 fL.

Statistical Analysis:

A comparison between the above-mentioned platelet parameters was performed on the data presented by mini tab program version 17, minimum, maximum, arithmetic mean and standard deviation, The significance of difference of different means (quantitative data) was tested using students t-test for difference between two independent means or one-way analysis of variance (ANOVA) test for difference among more than two independent means with Tukey's Pair wise comparisons were used. Means that do not share a letter are significantly different. The significance of difference of different percentages (qualitative data) was tested using Pearson Chi-square test (x²-test). Pearson correlation was calculated for the correlation between two quantitative variables with its t-test for testing the significance of correlation. To know the correlation between MAP and platelet (PLT) indices, pearson's correlation coefficient "r" was used which is a measure of strength and direction of linear dependence between two variables. The correlation coefficient value (r) either positive (direct correlation) or negative (inverse correlation), a statistical significance was considered whenever the P value was equal or less than 0.05.

Results

The results presented in this study were based on the analysis for a total data of 120 pregnant women in their third trimester. Sixty of them have preeclampsia (cases) in which 30 of them were with severe preeclampsia and 30 with mild preeclampsia and other 60 women were control group. The demographic and obstetric characteristics of all groups are shown in table (1). The mean maternal age ± SD was 24± 5.4 years in the preeclampsia and 26.5± 5.05 years in the control group, with very high significant (P = 0.009) difference between them. 41.67 % of cases

had positive history of previous preeclampsia with very high significant (P= 0.000) difference between preeclampsia and control group. There was no significant difference in the parity and gestational age with (P = 0.217 and P = 0.306), respectively, between cases and control group.

Table (1): Demographic and obstetric characteristics of preeclampsia and control groups

| Parameter (Range) | Cases "Preeclampsia" [n = 60] Mean ± SD | Control "Normotensive" [n = 60] Mean ± SD | P-value * |
|---|---|---|-----------|
| Maternal age (years) | 24± 5.4 | 26.5± 5.05 | 0.009 |
| Parity | 1.767 ±1.798 | 2.217 ±2.156 | 0.217 |
| Gestational age (weeks) | 38.1 ± 2.07 | 38.5 ± 2.06 | 0.306 |
| History of previous preeclampsia, no. (%) | 25 (41.67) | 0 (0.00) | <0.001 |

* *Independent T-test of two means was used for comparison in measurable variables whereas, chi-square test applied for categorical data.*

Table (2) Compared Platelet count PC (×10⁹/L), Mean platelet volume MPV (fL), platelet distribution width PDW (fL), PC/MPV distribution for the preeclampsia patients with control group. The present study found out the mean PC ± SD was 178.50 ± 41.80×10⁹/L in severe preeclampsia, 232.2 ± 63.6×10⁹/L for the mild preeclampsia group and 254.25 ± 41.08×10⁹/L in the control group. The mean ± SD for MPV was 11.807± 1.368 fl in severe preeclampsia, 10.663± 1.439 fL for the mild preeclampsia group and 9.472 ±0.945 fl in the control group. While mean ± SD for PDW was 20.390 ±2.415 fl in severe preeclampsia, 17.360 ±2.404 fL for the mild preeclampsia group and 14.443± 4.683 fl in the control group. Whereas mean PC/MPV ± SD was 16.81 ±5.69 in severe preeclampsia group, 23.00 ±8.63 for the mild preeclampsia group and 27.308 ±6.474 in the control group. The difference was statistically significant for all (P = <0.001).

Table (2): Comparison of platelet count and platelet indices among severe preeclampsia, mild preeclampsia, and control groups

| Parameters | Groups (Mean \pm SD) | | | P-value* |
|------------------------------------|------------------------|--------------------|--------------------|----------|
| | Severe preeclampsia | Mild preeclampsia | Normotensive | |
| No. of women | 30 | 30 | 60 | |
| Platelet count ($\times 10^9/L$) | 178.50 \pm 41.80 | 232.2 \pm 63.6 | 254.25 \pm 41.08 | <0.001 |
| Mean platelet volume (fl) | 11.807 \pm 1.368 | 10.663 \pm 1.439 | 9.472 \pm 0.945 | <0.001 |
| Platelet dist. W(fl) | 20.390 \pm 2.415 | 17.360 \pm 2.404 | 14.443 \pm 4.683 | <0.001 |
| PC/MPV | 16.81 \pm 5.69 | 23.00 \pm 8.63 | 27.308 \pm 6.474 | <0.001 |

* *One-way ANOVA-test with Tukey's Pair wise comparisons was used.*

Table (3) demonstrated the platelet count distribution for the preeclampsia and control groups. Twenty-one patients (70%) of severe Preeclampsia group with 8 patients (26.6%) in mild preeclampsia group and zero in control group had a platelet count between (150- 199) $\times 10^9/L$. While 6 patients (20%) of severe Preeclampsia group with 11 patients (36.6%) in mild group and 28 patients (46.6%) in control group had a platelet count between (200-249) $\times 10^9/L$. This was compared with 3 patients (10%) in severe preeclampsia group, 11 patients (36.6%) in mild preeclampsia group and 32 women (53.33%) in the control group had a platelet count of $\geq 250 \times 10^9/L$.

This difference between proportions for platelet count in the studied groups was statistically significant ($P = <0.001$).

Table (3): Distribution of platelets count distribution among the study groups

| Platelet count ($\times 10^9/L$) | Severe preeclampsia [n = 30] | | | | Normotensive [n = 60] | | P-value* |
|------------------------------------|------------------------------|----|-----|------|-----------------------|-------|----------|
| | No. | % | No. | % | No. | % | |
| | 150-199 | 21 | 70 | 8 | 26.6 | 0 | |
| 200-249 | 6 | 20 | 11 | 36.6 | 28 | 46.6 | <0.001 |
| ≥ 250 | 3 | 10 | 11 | 36.6 | 32 | 53.33 | |

* *Chi-square test was used, d.f = 4.*

Table (4) demonstrates the mean platelet volume MPV (fL) distribution for the preeclampsia and control groups. None in the severe preeclampsia group and mild preeclampsia group had MPV below 8fL. This compared with 4 women (6.67%) in the control group who had MPV below 8 fL. While one the patient in severe preeclampsia group (3.33%), 17 patients (56.67%) in mild preeclampsia group and 48 women (80%) in control group had MPV between 8 – 10.5 fL. However, 29 patients (96.67%) in severe preeclampsia group, 13 patients (43.33%) in the mild preeclampsia group, and 8 women (13.33%) in the control group had MPV equal or above 10.6 fL.

This difference between proportions for MPV in the studied groups was statistically significant ($P = <0.001$).

Table (4): Statistical difference of MPV among the study groups

| MPV (FL) | Severe preeclampsia [n = 30] | | Mild preeclampsia [n = 30] | | Normotensive [n = 60] | | P-value* |
|-------------|------------------------------|-------|----------------------------|-------|-----------------------|-------|----------|
| | No. | % | No. | % | No. | % | |
| < 8 | 0 | 0 | 0 | 0.00 | 4 | 6.67 | |
| 8 – 10.5 | 1 | 3.33 | 17 | 56.67 | 48 | 80 | <0.001 |
| ≥ 10.6 | 29 | 96.67 | 13 | 43.33 | 8 | 13.33 | |

* *Chi-squared test was used, d.f = 4.*

Table (5) listed the distribution of platelet width PDW (fL) for Preeclampsia and control groups. 6 patients (20%) in the severe preeclampsia group, 20 patients (66.67%) in the mild preeclampsia group,

and 40 women (66.67%) in the control group had PDW < 19.0 fL. This was compared with 24 patients (80%) in severe preeclampsia group, 10 patients (33.33%) in mild preeclampsia group and 20 women (33.33%) in control group had PDW ≥ 19.0 fL.

This difference between proportions for PDW in the presented study sample groups was statistically significant (P = <0.001).

Table (5): Distribution of platelet width (PDW) distribution between preeclampsia cases and control groups

| PDW (fl) | Severe preeclampsia [n = 30] | | Mild preeclampsia [n = 30] | | Normotensive [n = 60] | | P-value * |
|----------|------------------------------|----|----------------------------|-------|-----------------------|-------|------------------|
| | No. | % | No. | % | No. | % | |
| < 19.0 | 6 | 20 | 20 | 66.67 | 40 | 66.67 | <0.001 |
| ≥ 19.0 | 24 | 80 | 10 | 33.33 | 20 | 33.33 | |

* Chi-square test was used, d.f = 2.

To know the correlation between MAP and platelet (PLT) indices, Pearson’s correlation coefficient "r" was used which is a measure of strength and

direction of linear dependence between two variables.

It ranges from + 1 to -1, where 1 indicates total positive correlation, 0 indicates no association, and -1 indicate total negative correlation. Table (6) showed the correlation analysis between platelet counts (PC), Mean platelet volume (MPV) and platelet distribution width (PDW) parameters in preeclampsia groups with mean arterial pressure (MAP) there was a statistically significant inverse correlation between platelet count and the mean arterial pressure with r value of -0.517 and (P= <0.001). Also, compared the direct correlation between MPV, PDW and MAP with r values of 0.390, 0.411 and P values of (0.025, 0.001), respectively.

Table (6): Correlation between MAP, Platelet count and platelet indices in preeclampsia women, [n = 60]

| Platelet indices | Pearson’s correlation coefficient "r" | P-value |
|--------------------------------------|---------------------------------------|---------|
| MPA (mmHg) | | |
| Platelet count (×10 ⁹ /L) | -0.517 | <0.001 |
| MPV (fl) | 0.390 | 0.025 |
| PDW (fl) | 0.411 | 0.001 |

Discussion

In fact, there is a proof that in hypertension during pregnancy, the thrombocyte production time is noticeably reduced in comparison with normal pregnancy. Young platelets shown in a circulation are larger and represent a higher tendency to coagulation (18,21). It is obvious that endothelial injury led to increased platelet activation and thrombocyte consumption by creating micro thrombosis. In this condition, megakaryocytes in the bone marrow release platelets, which have a larger volume and more activity to compensate the thrombocyte consumption (22). Mean platelet volume MPV and platelet distribution width PDW are simply measured platelet parameters, which increase during thrombocyte activation. In order to get a larger surface platelet, change in shape during activation. Their shape changes from discoid to the spherical (12,23). The current study demonstrated significantly lower PC and PC to MPV ratio in

patients with severe preeclampsia compared with the mild preeclampsia and control groups This is agreed with Doğan et al. and AlSheeha et al.(24,25) who observed significantly lower PC and PC/MPV in preeclamptic women compared with the controls. The same studies found no significant difference in PDW in the preeclamptic women than healthy control groups. While in the current study, there was significant difference in MPV. AlSheeha et al.(25) study stated that there was no significant difference in MPV in the preeclamptic women compared with the controls but Doğan et al.(24) who documented a higher level of MPV in the preeclamptic women compared with the controls. The difference may probably be due to the different sample size included in their studies, different in equipment and/or method of the automatic blood count devices used to obtain hemogram. Current study showed that, platelet count decreased significantly with severe preeclampsia which is similar to the findings reported by Dogru et al.(13), Ustün et al.(26), Vijaya et al.(27), and Annam et al.(28) studies. In this study, in spite of the platelet count being normal in all the cases included, it was observed that platelet count decreased as the severity of the disease increased and was statistically significant when severe preeclampsia cases were compared with mild preeclampsia cases and control group.

Dhakre et al.(29) study revealed that, PDW and MPV increase as pregnancy progresses and these changes were more marked in the preeclampsia group complicated pregnancy than normotensive pregnancy while, the Platelet count decreases. Furthermore, Nooh et al.(30) and Dundar et al.(31) noted a significant increase MPV. Both of them also reported that, the increase MPV preceded onset of development of preeclampsia and started from 24 week of gestation. In both types of preeclampsia cases, platelet (PLT) count was statistically correlated with MAP in a negative manner, whereas MPV and PDW values were positively correlated with MAP with significant P value this was resembled the study done by Reddy et al.(32), While Unlü et al.(33) and Ceyhan et al.(34) observed no significant change between preeclamptic groups and controls in respect to relation between MAP and platelet count. It was suggested that the main cause for the inconsistency between these studies is probably due to the method of measurement of these platelet parameters, measurements performed in EDTA and change in a time-dependent manner. Also, different systems used in measurement can give different results, and this difference may extend up to 40% (29). Yang et al.(35) study showed that among three groups mild, severe PE and normotensive women the platelet count lowered in the preeclamptic group. However, PDW and the MPV increased. When compared with the levels of other platelet indices, the PDW showed significant elevation in the severe PE group. In the severe and mild preeclampsia groups, the PDW was statistically correlated with the mean arterial pressure (MAP) ($r = 0.231$; $P < 0.05$), this was corresponding with the results of the present study which was shown among platelet parameters, the PDW and MPV is significantly higher in women with severe preeclampsia than in women with mild preeclampsia and control group and is positively correlated with the MAP this was similarly reported by Vijaya et al.(27).

Conclusion:

Platelet parameters are cost effective, easily measurable, and practical markers can be used for monitoring of preeclampsia. Mean platelet volume (MPV) and platelet distribution width (PDW) has a significant linear correlation with mean arterial pressure in severe Preeclampsia and can be used as biomarker for severity of preeclampsia in high-risk pregnant women.

Recommendations

Ascertain the use of platelet parameters as an aid in the detection and management of a patient with preeclampsia.

Availability and accessibility of platelet parameters in the antenatal care health center will serve in discovering preeclampsia earlier especially the pregnant women who were at high risk for preeclampsia.

Authors Contributions:

The concept and study design, and the critical revision were done by Ahmed J. Alhusaynei, acquisition of data analysis, and the drafting of the manuscript was done by Hanan Dh. Nayyef.

Authors Declaration: Conflicts of interest: None.

We hereby confirm that all the Figures and Tables in the manuscript are ours. The project was approved by the Iraqi Board for Medical Specializations.

References

- AL-Maini EH, Ali SI. The correlation between serum levels of soluble platelet-selectin in preeclampsia with & without intrauterine growth retardation versus normal pregnancy. *Journal of the Faculty of Medicine Baghdad*. 2017 Apr 2;59(1):36-41. <https://doi.org/10.32007/jfacmedbagdad.591156>
- American College of Obstetricians and Gynecologists. *Hypertension in pregnancy. Report of the American College of Obstetricians and Gynecologists' task force on hypertension in pregnancy*. *Obstetrics and gynecology*, 2013 Nov; 122(5):1122-31.
- Payne B, Magee LA, von Dadelszen P. Assessment, surveillance and prognosis in preeclampsia. *Best practice & research Clinical obstetrics & gynaecology*, 2011 Aug 1; 25(4):449-62. <https://doi.org/10.1016/j.bpobgyn.2011.02.003>
- Edmonds K. *Hypertensive Disorders*. In: *Dewhurst's Textbook of Obstetrics & Gynaecology*. Ninth edition. UK: John Wiley & Sons Ltd. 2018 Nov 12:73-84.
- Abalos E, Cuesta C, Grosso AL, Chou D, Say L. Global and regional estimates of preeclampsia and eclampsia: A systematic review. *European Journal of Obstetrics & Gynecology and Reproductive Biology*, 2013 Sep 1; 170(1):1-7. <https://doi.org/10.1016/j.ejogrb.2013.05.005>
- Abass AE, Abdalla R, Omer I, Ahmed S, Khalid A, Elzein H. Evaluation of platelets count and indices in preeclampsia compared to normal pregnancies. *IOSR Journal of Dental and Medical Sciences*, 2016 Jul; 1(15):5-8. <https://doi.org/10.9790/0853-150750508>
- Kaur S, Khan S, Nigam A. Hematological profile and pregnancy: a review. *International Journal of Advances in Medicine*, 2014 Aug; 1(2):68-70. <https://doi.org/10.5455/2349-3933.ijam20140804>
- Mondal B, Paul DL, Sultana T, Rahman Q, Ahmed S, Fatema K et al. Assessment of platelet

- count and platelet indices in preeclampsia and eclampsia. *The American Journal of Innovative Research and Applied Sciences*, 2015; 1(3):80-4.
9. Asif N, Hassan K. Thrombocytopenia in pregnancy. *Hematology & Transfusion International Journal*, 2017; 5(5):1-4. <https://doi.org/10.15406/htij.2017.05.00133>
10. Bain BJ, Bates I, Laffan MA. Investigation of haemostasis. In: *Dacie and Lewis 46 Practical Haematology*. Twelfth edition. London: E-Book: Expert Consult: Online and Print. Elsevier Health Sciences. 2016 Aug 11: 336-409.
11. Jarjees SB. Evaluation of platelet count and mean platelet volume in normal and preeclamptic pregnant women. Iraq: Baghdad/college of medicine, 2011: 5. Dissertation for fellowship of the Arab Board for Medical Specialization in Obstetrics and Gynaecology.
12. Vagdatli E, Gounari E, Lazaridou E, Katsibourlia E, Tsikopoulou F, Labrianou I. Platelet distribution width: a simple, practical and specific marker of activation of coagulation. *Hippokratia*, 2010 Jan; 14(1):28.
13. Dođru HY, Yücel N, Pelit FÇ, Bolat G. The importance and evaluation of mean platelet volume on the severity of preeclampsia. *Perinatal Journal*, 2011; 19(3):108- 113. <https://doi.org/10.2399/prn.11.0193002>
14. Tzur T, Sheiner E. Is there an association between platelet count during the first trimester and preeclampsia or other obstetric complications later in pregnancy?. *Hypertension in pregnancy Taylor & Francis online*, 2013 Feb 1; 32(1):74-82. <https://doi.org/10.3109/10641955.2012.704109>
15. Wang F, Meng Z, Li S, Zhang Y, Wu H. Platelet distribution width levels can be a predictor in the diagnosis of persistent organ failure in acute pancreatitis. *Gastroenterology research and practice*, 2017 Dec 27:2017. <https://doi.org/10.1155/2017/8374215>
16. Boshnak N, Boshnaq M, Elgohary H. Evaluation of platelet indices and red cell distribution width as new biomarkers for the diagnosis of acute appendicitis. *Journal of Investigative Surgery*. 2018 Mar 4;31(2):121-9. <https://doi.org/10.1080/08941939.2017.1284964>
17. Tzur I, Barchel D, Izhakian S, Swarka M, Garach-Jehoshua O, Krutkina E, Plotnikov G, Gorelik O. Platelet distribution width: a novel prognostic marker in an internal medicine ward. *Journal of Community Hospital Internal Medicine Perspectives*. 2019 Nov 2;9(6):464-70. <https://doi.org/10.1080/20009666.2019.1688095>
18. Varol E, Akcay S, Icli A, Yucel H, Ozkan E, Erdogan D et al. Mean platelet volume in patients with prehypertension and hypertension. *Clinical hemorheology and microcirculation*, 2010 Jan 1; 45(1):67-72. <https://doi.org/10.3233/CH-2010-1327>
19. Thanoon AM, Jalal SD. Thrombocytopenia in Iraqi Pregnant Women. *Journal of the Faculty of Medicine Baghdad*. 2011 Jul 3;53(2):171-4. <https://doi.org/10.32007/jfacmedbagdad.532864>
20. NICE guideline criteria for preeclampsia diagnosis/2020. https://action-on-preeclampsia.org.uk/wp-content/uploads/2020/01/NICE_guidance_on-Preeclampsia_VB2020.pdf
21. Stekkinger E, Zandstra M, Peeters LL, Spaanderman ME. Early-onset preeclampsia and the prevalence of postpartum metabolic syndrome. *Obstetrics & Gynecology*, 2009 Nov 1; 114(5):1076-84. <https://doi.org/10.1097/AOG.0b013e3181b7b242>
22. Gabbe SG, Niebyl JR, Simpson JL, Landon MB, Galan HL, Jauniaux ER et al. *preeclampsia and hypertensive disorder*. In: *Obstetrics: Normal and Problem Pregnancies*. Seventh edition. London: E-book. Elsevier Health Sciences. 2016 Mar 18: 661-705.
23. A. Jagroop, I. Clatworthy, J. Lewin, DP Mikhailidis I. Shape change in human platelets: measurement with a channelyzer and visualisation by electron microscopy. *Platelets*, 2000 Jan 1; 11(1):28-32. <https://doi.org/10.1080/09537100075760>
24. Dođan K, Guraslan H, Senturk MB, Helvaciođlu C, İdil S, Ekin M. Can platelet count and platelet indices predict the risk and the prognosis of preeclampsia?. *Hypertension in pregnancy*, 2015 Oct 2; 34(4):434-42. <https://doi.org/10.3109/10641955.2015.1060244>
25. AlSheeha MA, Alaboudi RS, Alghasham MA, Iqbal J, Adam I. Platelet count and platelet indices in women with preeclampsia. *Vascular health and risk management*, 2016; 12:477. <https://doi.org/10.2147/VHRM.S120944>
26. Üstün YE, Dogan K, Türkçüođlu I, Üstün Y, Meydanl MM, Kafkasl A. Evaluation of hemoglobin and platelet levels in mild, moderate and severe preeclampsia. *Perinatal Journal*, 2007; 15(3): 93-8.
27. Vijaya C, Lekha MB, Shetty A, Geethamani V. Evaluation of platelet counts and platelet indices and their significant role in preeclampsia and eclampsia. *Journal of Evolution of Medical and Dental Sciences*, 2014 Mar 24; 3(12):3216-20. <https://doi.org/10.14260/jemds/2014/2269>
28. Annam V, Srinivasa K, Yatnatti SK, Med SD. Evaluation of platelet indices and platelet counts and their significance in preeclampsia and eclampsia. *International Journal of Biological & Medical Research*, 2011 Jan 31; 2(1):425-8.
29. Dhakre R, Nandmer GK, Sapkal R. Correlation of platelet indices with severity of preeclampsia: a prospective study from central India. *International Journal Reprod Contracept Obstet Gynecol*, 2018;

7(4):1416-20. <https://doi.org/10.18203/2320-1770.ijrcog20181327>

30. Nooh AM, Abdeldayem HM. Changes in platelet indices during pregnancy as potential markers for prediction of preeclampsia development. *Open Journal of Obstetrics and Gynecology*, 2015; 5(12):703. <https://doi.org/10.4236/ojog.2015.512099>

31. Dundar O, Yoruk P, Tutuncu L, Akyol Erikci A, Muhcu M, Ergur AR et al. Longitudinal study of platelet size changes in gestation and predictive power of elevated MPV in development of preeclampsia. *Prenatal Diagnosis: Affiliation With the International Society for Prenatal Diagnosis*, 2008 Nov; 28(11):1052-6. <https://doi.org/10.1002/pd.2126>

32. Reddy SG, Prasad CS. Significance of platelet indices as severity marker in nonthrombocytopenic preeclampsia cases. *Journal*

of laboratory physicians, 2019 Jul; 11(3):186. https://doi.org/10.4103/JLP.JLP_161_18

33. Ünlü BS, Acay A, Köken GG, Yıldız Y, Telciler E, Pektaş MK et al. The Role of Mean Platelet Volume in Preeclampsia. *Journal of Turgut Ozal Medical Center*, 2014 Sep 1; 21(3).

34. Ceyhan T, Beyan C, Başer İ, Kaptan K, Güngör S, Ifran A. The effect of preeclampsia on complete blood count, platelet count and mean platelet volume. *Annals of hematology*, 2006 May; 85(5):320-2. <https://doi.org/10.1007/s00277-006-0091-7>

35. Yang SW, Cho SH, Kwon HS, Sohn IS, Hwang HS. Significance of the platelet distribution width as a severity marker for the development of preeclampsia. *European Journal of Obstetrics & Gynecology and Reproductive Biology*, 2014 Apr 1; 175:107-11. <https://doi.org/10.1016/j.ejogrb.2013.12.036>

How to Cite this Article

Dhannoon Nayyef H, Jasim Alhusaynei A. Platelet Parameters in Nonthrombocytopenic Preeclampsia: A Case-Control Study. *JFacMedBagdad [Internet]. [cited 2023 Dec. 12];65(4). Available from: <https://iqjmc.uobaghdad.edu.iq/index.php/19JFacMedBaghdad36/article/view/2084>*

معلومات الصفائح الدموية لدى النساء الحوامل المصابات بمقدمات الارتجاج وبدون نقص في عدد الصفائح الدموية: دراسة مقارنة

د. احمد جاسم الحسيني /م. الخنساء التعليمي للولادة / الموصل
د. جنان دو النون نايف / م. الخنساء التعليمي للولادة/ الموصل
الخلاصة

الخلفية: مقدمات الارتجاج هو اضطراب خاص بالحمل متعدد الأجهزة، يتميز بتطور ارتفاع ضغط الدم والبييلة البروتينية التي تنشأ بعد الأسبوع العشرين من الحمل في امرأة كانت ذات ضغط طبيعي سابقاً ويرتبط الحمل بالتكيف الفسيولوجي الذي يؤدي الى تغيير في نظام الدم ومن ضمنها معلومات الصفائح الدموية.

هدف الدراسة: دراسة لمقارنة العلاقة بين عدد الصفائح الدموية (pc) ومؤشرات الصفائح الدموية وهي متوسط حجم الصفائح الدموية (MPV) وتوزيع الصفائح الدموية (PDW) ونسبة عدد الصفائح الى حجمها في النساء المصابات بمقدمات الارتجاج مع النساء ذوات الحمل الطبيعي. **المرضى والطريقة:** تم إجراء دراسة الحالة والشاهد في قسم النسائية والتوليد في مستشفى الخنساء التعليمي للأطفال والولادة، الموصل، العراق على مدى فترة تمتد من الأول من شباط ٢٠٢٠ إلى نهاية كانون الأول ٢٠٢٠. شاركت عينة من ١٢٠ امرأة حامل في الدراسة ممن استوفين معايير الاشتمال. ٦٠ منهن تم تشخيص إصابتهن بمقدمات الارتجاج (٣٠ امرأة حامل مع مقدمات الارتجاج الخفيف و ٣٠ امرأة حامل مع مقدمات الارتجاج الشديد و ٦٠ امرأة حامل حملاً طبيعياً كمجموعة شاهد. تم قياس عدد الصفائح الدموية ومتوسط حجم الصفائح الدموية كجزء من تعداد الدم الكامل.

النتائج: علاقة خطية معنوية بين متوسط حجم الصفائح الدموية (MPV) وعرض توزيع الصفائح الدموية (PDW) بمتوسط ضغط شرياني في مجموعة تسهم الحمل بقيمة p (0.025 و 0.001) على التوالي وعلاقة عكسية معنوية بين عدد الصفائح الدموية ومتوسط الضغط الشرياني مع p تمت ملاحظة قيمة (>0.001).

الخلاصة: تعتبر مؤشرات الصفائح الدموية فعالة من حيث التكلفة، وسهلة القياس، وعلامات عملية لرصد مقدمات الارتجاج، وان متوسط حجم الصفائح الدموية وعرض توزيعها له علاقة طردية كبيرة مع متوسط الضغط الشرياني في حالة مقدمات الارتجاج الشديد. **الكلمات المفتاحية:** مقدمات الاتعاج، معلومات الصفائح الدموية، عدد الصفائح الدموية، حجم الصفائح الدموية، توزيع الصفائح الدموية.