

Association between Body Mass Index and Depressive Symptoms among Adolescent Females in Baghdad, AlKarkh during 2022

Masarrah M. Alkabban^{1*}  , Lujain A. Alkhazrajy²  

¹Baghdad AL-Karkh Health Directorate, MOH, Baghdad, Iraq.

²Department of Family & Community Med. Al-Kindy College of Medicine- University of Baghdad, Baghdad, Iraq.



©2024 The Author(s). Published by College of Medicine, University of Baghdad. This open-access article is distributed under the terms of the Creative Commons Attribution 4.0 International License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract:

Background: Due to their high prevalence rates, adolescent obesity and depression are among the most renowned public health concerns globally. Obesity is considered the leading cause of morbidity and death in Iraq in 2019. There is scarce research and surveys regarding the relation between BMI and depression especially in young Iraqi females.

Objectives: To explore the level of depression among adolescent girls, and to determine its association with Body Mass Index in the studies group.

Methods: Beck's depression inventory questionnaire was used in 11 high schools for girls that were selected randomly from the schools in Baghdad/ AlKarkh from 1st of January 2023 till end of May 2023. A sample of 350 adolescent girls was included in the study. The height and weight of each participant was measured and BMI was calculated. Tables and graphs were constructed to describe the data. The Chi-square test was used to test the association between categorical data. A P value of less than 0.05 was considered as statistically significant.

Results: Of the studied group, 14% were underweight, 59% normal weight, 21% overweight and 6% obese. As for mood status, 16% had mild mood disturbance, 24.3% had borderline clinical depression, 13.4% had moderate depression and 3.1% had severe depression. Severe depressive symptoms were found in 15% of obese girls, significantly higher than that among overweight (1.4%), normal weight (2.4%) and underweight (3.4%).

Conclusions: There was a significant association between obesity and severe depression, and a positive significant correlation between BMI and depression.

Keywords: Adolescents; BMI; Depressive symptoms; Iraqi girls; Obesity.

Received: April, 2023

Revised: May 2023

Accepted: June, 2023

Published: Jan., 2024

Introduction:

Many people feel sad at times, but these feelings usually disappear within a couple of days. When someone has a depressive disorder, it impedes daily life and normal work and affects a person's conduct with family and friends (1). Depression is defined as "a mood state categorized by a sense of inadequacy, a feeling of hopelessness, a decrease in activity or reactivity, negativity, sadness and associated symptoms. It is one of the most common mental health disorders, which is also considered the foremost cause of ill health worldwide (2). Risk factors for depression are female sex, low educational level, low socioeconomic status, comorbid anxiety disorders, family history of depression, personal history of depression, comorbid physical chronic diseases, and exposure to stressful life events (3). It is believed that 5% of adults worldwide experience depression, which can cause a wide range of mental and physical issues and reduce the capacity to function both at work and at home (4).

According to the 2021 World Health Organization

(WHO) report, nearly 322 million people (4.4%) worldwide live with depression (5). According to a recent Iraqi Ministry of Health (MOH) report, 16.5% of Iraqi people older than 15 years of age suffer from psychological diseases.

Depression affects nearly 10.9% of individuals aged 18-25 years (6). Depression can lead to suicide which is the second leading cause of death in young adults (15-29 years old) globally (7).

Obesity is considered one of the non-communicable diseases (NCDs) which were the leading cause of morbidity and death in Iraq in 2019 (8). Overweight and obesity are defined as abnormal or excessive fat accumulation that may impair health (9). Adolescence is the phase of life between childhood and adulthood, from ages 10 to 19. It is a unique stage of human development and an important time for laying the foundations of good health (10). Body image is made in childhood from the interaction with parents, siblings, friends, and schoolmates. Hence, whether these interactions were positive or negative will influence on the individual's self-esteem (11). Even though there are many studies on the

* Correspondence author: alkabbanmasarrah@gmail.com

relationship between obesity and depression, this topic was selected because there is scarce research work regarding the relationship between BMI and depression especially in young Iraqi females.

Subjects and Methods:

This Cross-sectional study was conducted in Baghdad in 11 high schools chosen, from Baghdad/ AlKarkh (First and Second Directorates of Education) from 1st of January 2022 till end of May 2023. A multistage random sampling method was used to select a sample of 350 female adolescents (10-12 students were chosen by simple random sampling from each class, from each school).

Exclusion criteria: Adolescents who had experienced divorce of parents, remarriage of parents, death in the family, or who have any acute or chronic physical and psychiatric disease.

The data was collected by using the (Beck's depression inventory questionnaire) that was used in previous studies (12), which included 21 questions designed to measure depressive symptoms. The questionnaire was validated by three senior physicians (community medicine, family medicine and a psychiatrist). The researcher visited all the selected schools and a direct interview was made with each selected student, to fill the questionnaire. The height and weight of each student was measured using a weight and height meter (Seca). The BMI was calculated. The girls were grouped according to their BMI to: Underweight, normal weight, overweight and obese. (13)

Approval was obtained from the administrations of the selected schools before starting the data collection. Verbal consent was obtained from each selected student after full clarification of the aims of the study and assuring the confidentiality of the collected data.

The data were loaded into SPSS V26 statistical programs. Tables and graphs were used for descriptive purposes. The Chi-square test was used to test associations between categorical data. The correlation coefficient was calculated to test its significance between numerical variables. A P value of less than 0.05 was considered significant.

Results:

Figure (1) show that 40.3% of the participants were 16 years old, 41.1% were 17 years old, 16% were 18 years old and 2.6% were 19 years old.




Figure 1: Age distribution of the studied sample

The study also shows that 14% were underweight, 59% were normal weight, 21% overweight and 6% were obese, figure (2).




Figure (2): BMI distribution of the studied sample

While figure (3) shows that 43.1% of studied students had no depressive symptoms, 16% had mild mood disturbance, 24.3% had borderline clinical depression, 13.4% had moderate and 3.1% had severe depression.




Figure (3): Distribution of studied sample according to their depressive symptoms

Table 1 shows that 15% of obese students had severe depressive symptoms which was significantly higher than the rate of severe depressive symptoms among overweight (1.4%), normal weight (2.4%) and underweight (3.4%), P = 0.003.

Table (1): Association between weight status and depressive symptoms

Weight status	Normal		Mild mood disturbance		Borderline clinical depression		Moderate depression		Severe depression		
	N	%	N	%	N	%	N	%	N	%	
Under weight	22	43.1	7	13.7	8	15.7	12	23.5	2	3.9	51
Normal weight	95	46.1	29	14.1	57	27.7	20	9.7	5	2.4	206
Overweight	32	43.8	16	21.9	13	17.8	11	15.1	1	1.4	73
Obese	2	10.0	4	20.0	7	35.0	4	20.0	3	15.0	20
Total	151	143	56	69.7	85	96.2	47	68.3	11	22.7	350
<i>p</i> value	0.003										

Table 2 shows that there is no significant association between the age of the students and depressive symptoms, $p = 0.195$.

Table (2): Association between age of participants and depressive symptoms

Age (Years)	Normal		Mild mood disturbance		Borderline clinical depression		Moderate depression		Severe depression		Total
	N	%	N	%	N	%	N	%	N	%	
16	60	42.6	25	17.7	33	23.4	18	12.8	5	3.5	141
17	64	44.4	22	15.3	40	27.8	13	9.0	5	3.5	144
18-19 *	27	41.5	9	13.8	12	18.5	16	24.6	1	1.5	65
Total	151	143	56	69.7	85	96.2	47	68.3	11	22.7	350
<i>P</i> value	0.195										

because of small number of cases in age 18 and 19 year and for suitable calculation of the chi-square test, cases of these 2 ages were merged in one group

A weakly significant positive correlation was noticed between BMI and depressive symptoms score ($R = 0.111$, $P = 0.038$), (Fig 4).




Figure (4): Correlation between BMI and depressive symptoms score

Discussion:

Due to their high prevalence, morbidity, and death rates, adolescent obesity and depression are among the most renowned public health concerns globally (14). Therefore, a substantial amount of research has focused on the factors that contribute to the development of obesity and depression.

Results similar to those of our study regarding body weight of adolescent girls were reported by a study from Iran in 2016, where 59.8% of the study sample were of normal weight, 12% overweight and 7.3% obese (15). In a Korean study on adolescents, 58.9% were of normal body weight, 17.0% were overweight and obese and 24.1% were underweight (16).

As for depressive symptoms, the above study from Iran showed that 27.3% were normal with no depressive symptoms, 40.5% had minor depressive symptoms, 19.5% had moderate depression, and 12.8% had major depression (15).

An examination of a cohort of obese patients in the United Kingdom regarding the influence of BMI on depression revealed a stepwise pattern of increasing risk of depression among those with a BMI higher than 30 kg/m², with the risk doubling among those with a BMI greater than 60 kg/m² (17). In a study published in 2013 in USA, greater BMI levels were related to depressive symptoms completely through a psychosocial factor; the negative body image (18). The 2016 study from Iran showed that there was a positive and significant connection between BMI and depression among high school female students (15). A report on the search of three major databases in 2016 showed that overweight and obesity were adversely related to a number of psychiatric comorbidities, including depression (19). A systematic review on the subject in 2017 showed that the majority of cross-sectional researches indicated a strong relationship between obesity and depression (20). Small and Ablasca (2016) acknowledged the ambiguous relationship between obesity and depression. This study revealed a highly significant association between body weight and depressive symptoms, as obese girls report a greater prevalence of severe depression (21). All the above-mentioned studies and reviews are consistent with the results of the current study.

A study from Turkey in 2020 showed no statistically significant association between depression symptoms and the age of the participants (22) similar to our findings. Other variables may influence the relationship between obesity, anxiety, and depression. Low physical activity, an unhealthy diet, and sleep disturbances are some of the confounders for both depression and obesity (23, 24). In addition, obese adolescents are often harassed or bullied

because of their weight, which may contribute to anxiety and depression symptoms (25).

Obesity is connected with subclinical systemic inflammation and oxidative stress, which are identified as key etiological components of depression (23). Furthermore, obesity is strongly influenced by genetic factors, with an estimated heritability of 60% regarding BMI. Genetic susceptibility to the common form of obesity appears to be polygenic (27).

Regarding the comparison to Iraqi literature, in 2020, females with a healthy nutritional status were less likely to be depressed, whereas a larger consumption of fatty and unhealthy meals was connected with increased anxiety (28-30).

Conclusions: There is a significant association between obesity and depression. A positive and significant correlation between BMI and depression was also noticed.

Author Declaration:

Conflicts of Interest: None.

We hereby confirm that all the Figures and Tables in the manuscript are mine/ ours. Besides, the Figures and images, which are not mine /ours, have been given permission for re-publication and attached to the manuscript. Authors sign on ethical consideration's approval-Ethical Clearance: The project was approved by the local ethical committee in the Ministry of Health.

Authors' contributions:

Study conception & design: (Lujain A. Alkhazrajy). Literature search: (Masarrah M. Alkabban). Data acquisition: (Masarrah M. Alkabban). Data analysis & interpretation: (Lujain A. Alkhazrajy). Manuscript preparation: (Masarrah M. Alkabban). Manuscript editing & review: (Lujain A. Alkhazrajy).

References

1. Nigatu, Y. T. *Obesity and Depression: An Intertwined Public Health Challenge. [Thesis fully internal (DIV), University of Groningen]. University of Groningen 2015.*
2. Banka P: *The Relationship Between Childhood Depression, Obesity and Poor School Performance among Urban Disadvantaged Children (Thesis). Trinity College Dublin. School of Nursing & Midwifery. General Nursing, Trinity's Access to Research Archives. 2018:1-5*
3. Darwish L, Beroncal E, Sison MV, and Swardfager W: *Depression and health status in type 2 diabetes mellitus general perspectives. Diabetes, Metabolic syndrome and Obesity Journal vo. 11 2018, p11-12* <https://doi.org/10.2147/DMSO.S106797>.
4. World Health Organization Health Topics / Depression. https://www.who.int/health-topics/depression#tab=tab_1/2022.
5. World Health Organization. *Depression and Other Common Mental Disorders: Global Health Estimates.*

<http://apps.who.int/iris/bitstream/10665/254610/1/WHO-MSD-MER-2021.2-eng.pdf?ua=1>.

6. World Health Organization Department of Mental Health and Substance Abuse. *Depression: A global public health concern.*

https://www.who.int/mental_health/management/depression/who_paper_depression_wfmh_2012.pdf.

7. World Health Organization 2023. *Depression.*

<https://www.who.int/news-room/fact-sheets/detail/depression>.

8. World Health Organization Iraq: *Non-communicable Diseases (NCD) Country Profiles. WHO2022 Available online:*

<https://www.emro.who.int/iraq/priority-areas/noncommunicable-diseases.html>.

9. World Health Organization. *Obesity and overweight* <https://www.who.int/news-room/fact-sheets/detail/obesity-and-overweight/2022>.

10. World Health Organization / Adolescent health / https://www.who.int/health-topics/adolescent-health#tab=tab_1%202022.

11. Peña YO, Carvajal AS, Luna MO, Bojórquez RC. *Scientific Research. Gender, and Satisfaction of Body Image in High School Students of Yucatan, Mexico. Autonomous University of Yucatan, Mérida, Mexico. 2019 10(1)* <https://doi.org/10.4236/psych.2019.101003>.

12. Beck depression inventory, second edition, Czech version: *Demographic correlates, factor structure and comparison with foreign data. International Journal of Psychiatry in Clinical Practice, https://www.tandfonline.com/journals/ijpc20, 10 Oct 2022* 19. CDC (centres for Disease Control and Prevention)/<https://www.cdc.gov/obesity/data/adult.html> May 17, 2022.

13. World Health Organization. *A healthy lifestyle - WHO recommendations*

<https://www.who.int/europe/news-room/fact>

14. Blasco BV, García-Jiménez J, Bodoano I, Gutiérrez-Rojas L. *Obesity and Depression: Its Prevalence and Influence as a Prognostic Factor: A Systematic Review. Psychiatry Investig. 2020; 17:715-724.* <https://doi.org/10.30773/pi.2020.0099>.

15. Tashakori A, Riahi F, and Mohammadpour A. *The Relationship between Body Mass Index and Depression among High School girls in Ahvaz. Advances in Medicine. 2016: 1*

<https://doi.org/10.1155/2016/3645493>.

16. Lim Y and Kim B. *Body Mass Index, Body Weight Perception, and Depressed Mood in Korean Adolescents. J Korean Acad Child Adolesc Psychiatry 2017; 28(1): 31-37* <https://doi.org/10.5765/jkacap.2017.28.1.31>.

17. Moussa OM, Ardissino M, Kulatilake P, Faraj A, Muttoni E, Darzi A, et al. *Effect of body mass index on depression in a UK cohort of 363 037 obese patients: A longitudinal analysis of transition. Clin Obes. 2019;9(3):e12305.*

<https://doi.org/10.1111/cob.12305>.

18. Eidsdottir ST, Kristjansson AL, Sigfusdottir IG, Garber CE, Allegrante JP. *Association between higher BMI and depressive symptoms in Icelandic adolescents: the meditational function of body image.*

Eur J Public Health 2014 Dec;24(6):888-92
<https://doi.org/10.1093/eurpub/ckt180>.

19. Rankin J, Matthews L, Cogley S, Han A, Sanders R, Wiltshire HD, et al. Psychological consequences of childhood obesity: Psychiatric comorbidity and prevention. *Adolesc. Health Med. Ther.* 2016; 7: 125.
<https://doi.org/10.2147/AHMT.S101631>.

20. Mühlig Y, Antel J, Föcker M, Hebebrand J. Are bidirectional associations of obesity and depression already apparent in childhood and adolescence as based on high-quality studies? A systematic review. *Obes. Rev.* 2016;17:235-249.
<https://doi.org/10.1111/obr.12357>.

21. Small L, Aplasca A. Child obesity and mental health: A complex interaction. *Child Adolesc. Psychiatr. Clin. N. Am.* 2016; 25: 269-282.
<https://doi.org/10.1016/j.chc.2015.11.008>.

22. Sagar R, Gupta T. Psychological aspects of obesity in children and adolescents. *Indian J. Pediatr.* 2018;85:554-559. <https://doi.org/10.1007/s12098-017-2539-2>.

23. Berk M, Williams LJ, Jacka FN, O'Neil A, Pasco JA, Moylan S, et al. So depression is an inflammatory disease, but where does the inflammation come from? *BMC Med.* 2013; 11:200.
<https://doi.org/10.1186/1741-7015-11-200>.

24. Weisz JR, Chorpita BF, Palinkas LA, Schoenwald SK, Miranda J, Bearman SK, et al. Testing standard and modular designs for psychotherapy treating depression, anxiety, and conduct problems in youth: a randomized effectiveness trial. *Arch Gen Psychiatry.* 2012;69(3):274-82.
<https://doi.org/10.1001/archgenpsychiatry.2011.147>.

25. Puhl RM, King KM. Weight discrimination and bullying. *Best Pract Res Clin Endocrinol Metab.* 2013;27(2):117-27.
<https://doi.org/10.1016/j.beem.2012.12.002>

26. Alkhazrajy LA. Relationship between screen watching and overweight or obesity in a Sample of Iraqi adolescents. *AL-Kindy College Medical Journal.* 2013 Jan 30;9(1):80-8.
<https://jkmc.uobaghdad.edu.iq/index.php/MEDICAL/article/view/558>

27. Ramadan S, Abd Al Kareem F, Alkhazrajy LA. Genetic Polymorphisms in the Fat Mass and Obesity Associated Gene Confers Risk of Obesity in Iraqi Population. *American Journal of Infectious Diseases* 7(2)40-44, <https://doi.org/10.3844/ajidsp.2011.40.44>

28. Alkhazrajy LA, Abdulla R, Baher HH. The association between quality of habitual diet and mental health status among Iraqi women attending primary health care centers. *Archivos Venezolanos de Farmacologia y Terapeutica.* 2020; 39 (5): 604-609. <https://doi.org/10.5281/zenodo.4264720>.

29. Al Maliki A, AL Lami F, Al Aboudi S. Prevalence and Determinants of Depression among Diabetic Patients, Babel Province, Iraq, 2013-2014. *JfacMedBagdad.* 2015; 56(4):411-6.
<https://doi.org/10.32007/jfacmedbagdad.564558>.

30. Jasim HM, Ali BM, Al-Kaseer EA, AL-Diwan J. Depression among Females in AL-Sader city, Baghdad, Iraq. *JFacMedBagdad [Internet].* 2017 Oct. 1 [cited 2023 Apr. 1]; 59(3):231-3.
<https://doi.org/10.32007/jfacmedbagdad.59394>.

How to Cite

Alkabban MM, Alkhazrajy LA. Association between Body Mass Index and Depressive Symptoms among Adolescent Females in Baghdad, AlKarkh during 2022. *J Fac Med Baghdad.* 2024;65(4).

Available from:

<https://iqjmc.uobaghdad.edu.iq/index.php/19JFacMedBaghdad36/article>

العلاقة بين مؤشر كتلة الجسم وأعراض الاكتئاب لدى الفتيات في بغداد-الكرخ خلال عام 2022

مسرة مزاحم¹ ، لجين انور الخزرجي²

¹دائرة صحة بغداد الكرخ، وزارة الصحة، بغداد، العراق

²فرع طب الاسرة، كلية طب الكندي، جامعة بغداد، بغداد، العراق

الخلاصة

الخلفية: نظراً لارتفاع معدلات الإصابة والوفاة، تعد سمنة المراهقين والإكتئاب من أكثر إهتمامات الصحة العامة شهرة على مستوى العالم تعتبر السمنة السبب الرئيسي للاعتلال والوفاة في العراق في عام 2019. هناك أبحاث واستطلاعات شحيحة بخصوص العلاقة بين مؤشر كتلة الجسم والاكتئاب خاصة لدى الشباب العراقيين.

هدف الدراسة: فحص مستوى الإكتئاب لدى الفتيات المراهقات وتحديد العلاقة بين الإكتئاب ومؤشر كتلة الجسم لدى المراهقات في عينة الدراسة. **طرق البحث:** عن طريق استبيان بيك للإكتئاب الذي أجري في بغداد في 11 مدرسة ثانوية تم اختيارها عشوائياً من بغداد / الكرخ وتم طرح الاستبيان ل 350 مراهقة تم اختيارهن بطريقة عشوائية من المدارس للفترة من 1 كانون الثاني 2023 والى نهاية ايار 2023 وبعدها تم قياس طول ووزن كل طالبة باستخدام مقياس خاص للوزن والطول وتم حساب مؤشر كتلة الجسم لكل طالبة وتم تقسيم الطالبات المشمولات وفقاً لوزنهن. تم استخدام اختبار مربع كاي لمعرفة اهمية الارتباط بين المتغيرات واعتبرت قيمة $P < 0.05$ كنقطة لتميز الاحتمالية.

النتائج: 14% من الخاضعات للدراسة كن يعانين من نقص الوزن، 59% وزن طبيعي، 21% وزن زائد و 6% سمنة. من الطالبات اللاتي خضعن للدراسة لم يكن لديهن أعراض اكتئاب 43.1%، 16% لديهن اضطراب مزاجي خفيف، و 24.3% كن على حدود الإكتئاب السريري، و 13.4% كان لديهن اكتئاب متوسط بينما الإكتئاب الشديد وجد في 3.1% من الحالات. 15% من الطالبات اللواتي يعانين من السمنة المفرطة كانت لديهن أعراض إكتئاب حادة وهي أعلى بكثير من معدل أعراض الإكتئاب الحادة بين اللواتي يعانين من زيادة في الوزن (1.4%) وأيضاً أكثر من ذوات الوزن الطبيعي (2.4%) واللواتي يعانين من نقص الوزن (3.4%).

الإستنتاج: هناك إرتباط كبير بين السمنة والإكتئاب الشديد وايضاً إرتباط كبير بين معدل كتلة الجسم والإكتئاب.

الكلمات المفتاحية: المراهقون، مؤشر كتلة الجسم، السمنة، أعراض الإكتئاب، الفتيات العراقيات