

Antioxidant Effects of Selenium Nanoparticles Prepared from *Eruca Sativa* Extract on Ketoconazole –Induced Testicular Oxidative Damage in Male Rats

DOI: <https://doi.org/10.32007/jfacmedbagdad.6612174>

Basman Q. Shareef^{1*}  Huda I. AlQadhi¹  Shayma'a A. Jamal² 

¹Department of Pharmacology, College of Medicine, Baghdad University, Baghdad, Iraq.

²Department of Anatomy, College of Medicine, Baghdad University, Baghdad, Iraq.



©2024 The Author(s). Published by College of Medicine, University of Baghdad. This open-access article is distributed under the terms of the Creative Commons Attribution 4.0 International License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract:

Background: Recent studies have shown that the anti-fertility effects of Ketoconazole can be minimized by taking an antioxidant nanoparticles based-plant extract in combination with the medication or by taking it after medication. One of the effective methods to improve its medicinal properties and decrease the toxicities of Selenium is by formulating it as nanoparticles. However, this is still a challenging strategy.

Objectives: The aim of this study was to analyze the impact of nano selenium, which was prepared from *Eruca Sativa* extract, on testicular oxidative stress parameters in rats that were treated with ketoconazole. The study will focus on the evaluation of glutathione peroxidase, superoxide dismutase, and malondialdehyde as indicators of oxidative stress.

Methods: In this study, a 1% w/v solution of plant extract was added to a solution of 10mM sodium selenite in different ratios, and placed on a magnetic stirrer in the dark for 12 hours at a temperature of 50C° and pH9. The solution was then left for 48 hours, and the optimal fabricated selenium nanoparticles were selected for further characterization. Forty-eight rats were divided into six groups with eight animals in each. Group A was the negative control, while Group B was given oral Ketoconazole at a dose of 50mg/kg for fourteen days. Group BC, BD, BE1, and BE2 were given oral Ketoconazole at a dose of 50 mg/kg for fourteen days, followed by 200mg/kg *Eruca Sativa*, 0.5mg/kg oral sodium selenite, 0.25mg/kg oral nano selenium and 0.5mg/kg oral nano selenium for 28 days, respectively. Finally, the animals were euthanized and their testicle anti-oxidant parameters were evaluated.

Results: Significant increases in glutathione peroxidase and superoxide dismutase (p -value < 0.001), and a decrease in malondialdehyde levels (p -value < 0.001) were observed in groups treated with nanoparticles compared to the control group.

Conclusion: Nano forms prepared by *Eruca sativa* extract exhibit significant antioxidant effects on testicular tissues while being available for several metabolic, biological, and physiological functions.

Keywords: Glutathione Peroxidase; Malondialdehyde; Nano Selenium; Oxidative Stress; Selenium; Superoxide Dismutase.

J Fac Med Baghdad
2024; Vol.66, No. 1
Received: Jun., 2023
Accepted: Nov 2023
Published: Aprl.2024

Introduction

Selenium is one of the most significant trace elements in medicine and biology. It is found as incorporated Selenoproteins rich in Selenocysteine. It acts as a co-factor for various enzymes involved in oxidoreductase activities, such as glutathione peroxidase and thioredoxin reductase (1). These enzymes possess a cytoprotective effect, and their activation helps boost fertility, reduce inflammatory reactions, and provide a chemo-preventive effect against different types of cancers (2). In this regard, free radical-induced oxidative stress is thought to be the most important risk factor that threatens testicular functions; therefore, an antioxidant defense system which is represented by enzymes

incorporated with selenoproteins can counteract these free radicals prevent their formation in the testicular cells (3, 4).

Nanotechnology is a novel growing field with important applications in science and technology. In the pharmaceutical industry, nanoparticles act as carrier platforms to transport drugs to their target sites of action. It is based on the creation of small particles at the submicroscopic level (<100 nm) (5), which serves to enhance biological activity, minimize toxicity, and allow for the regulated release of medications, especially in capsule form (6).

Metal nanoparticles can be prepared by using either chemical or biological reducing agents. Implementing plant extracts with prominent antioxidant effects may help achieve that target.

* Corresponding Author:

basmaan.qasem1106b@comed.uobaghdad.edu.iq

Such formulations may achieve two targets using one stone, as the nanoparticle formulation will enhance the permeation of both Selenium and the plant extract into that target site, and this may help achieve a synergistic effect (7).

Ketoconazole is an antifungal drug that was reported to reduce epididymis sperm concentration, decrease the weight of male reproductive organs, especially the testes, and decrease serum testosterone levels (8), in addition to other pathological changes characterized by atrophied seminiferous tubules, malformed germ cells, and degenerated Sertoli cells. Interestingly, co-administration or post-treatment with antioxidant phytochemicals of plant extract could prevent such testicular damage (9).

This study was designed to investigate changes in testicular oxidative stress parameters (Superoxide dismutase "SOD", Malondialdehyde "MDA" and Glutathione peroxidase "GPX") in response to antioxidant effects of nano selenium prepared from *Eruca Sativa* extract in male rats.

Materials and Methods

Eruca Sativa Extract: Preparation and Characterization

One hundred grams of dried leaves powder of *Eruca sativa* was macerated with 2 liters n-hexane (99.8%) for four days for defatting plant material, then 80% ethanol was added, and the solution was filtered and evaporated using a rotary evaporator (4 rpm at 40°C). Finally, ethyl acetate and sodium Disulphate were added and the solution was then filtered, left to dry, and stored in a dark, sterile screw bottle at 4°C until used. The extract was characterized by measuring the total flavonoid content and its antioxidant power using a reducing power assay (10).

A. Measurement of Total Flavonoid Content

Flavonoids were estimated using a technique described by Ibraheem *et al*, 2014, where each mg of Gallic acid equivalent per gram of extract had its absorbance measured against a blank at 510 nm, and a calibration curve was analyzed (11).

B. Measurement of the Extract's Total Antioxidant Activity (Reducing Power Assay)

The extract's reducing power was measured according to the method prescribed by Ibraheem *et al*, 2012; the absorbance of different concentrations of the extract was measured at 700 nm and compared with the control (DW) (12). The percentage of absorption increase was sketched with different concentrations of Vitamin C as a standard (13).

Preparation of Selenium Nano Composite Using *Eruca Sativa* Extract

Briefly, a solution of 1% w/v of the extract was loaded to a solution of 10 mM sodium selenite at the four proportions (1:2, 1:4, 1:10, and 1:20 sodium selenite/ extract). The former was prepared by dissolving 0.172 gm. of sodium selenite (Na₂SeO₃) in 100 ml of deionized water. Then, the final mixture

was mixed using a hot plate magnetic stirrer (temp. 50 °C & pH of around 9) in the dark for 12 hrs. It was then autoclaved (temp. 121°C & gas pressures of 1.5 bar) for 15 min (14). Next, the autoclaved mixture was exposed to filter sterilization using a Millipore filter membrane (0.22µm) and ultra-sonic vibration (20 KHz by Q700 sonicator) to ensure an even dispersion of the nanoparticles in the liquid and to get micron-sized colloidal particles. Finally, the prepared colloid was sent for the characterization processes (15).

Characterization of the Nanocomposite

A. UV-Vis spectra analysis

B. Fourier transform infrared spectroscopy (FTIR).

C. Dynamic Light Scattering (DLS)

The samples were analyzed at the University of Technology's Nanotechnology Department.

Experimental design

A. Animals and ethical issues

This study was carried out on forty-eight healthy male adult albino rats (weighing 301± 24.28 grams and aged 10±2 weeks). They were gathered from the animal house at AL-Nahrin University/ College of Pharmacy and the Iraqi Center for Cancer and Genetic Research at AL-Mustansiriyah University. The study was approved by the Institutional Review Board (IRB)/College of Medicine /Baghdad University (Pharma Comed uvB 23.6) all the procedures and experiments were performed following the rules and legitimacy of the animal ethics committee /College of Pharmacy/ Al-Nahrin University.

Animals grouping: The experiment was implemented from the first of March 2022 to the end of August 2022. First, the animals were randomly divided into six groups with eight rats in each; Group A: -ve control, orally administered with 2 ml distilled water (DW) daily via gavage tube for 42 days. Group B: +ve control, orally received 2 ml of Ketoconazole (KET) solution (50mg/kg /day) via gavage tube (16) for 14 days, followed by 2 ml distilled water for 28 days. Group BC: received 2 ml of ketoconazole solution (50 mg/kg/day) orally for 14 days, followed by a therapeutic dose of *E. Sativa* extract (200mg/kg/day) orally (17) via gavage tube once daily for 28 days. Group BD: received 2 ml of ketoconazole solution (50mg/kg/day) orally for 14 days, followed by a therapeutic dose of sodium selenite solution (0.5 mg/kg/day) orally (18) via gavage tube once daily for 28 days.

Group BE: Nano Selenium group, which was subdivided into: Group BE₁ and BE₂: received 2 ml of ketoconazole solution (50 mg/kg/day) orally for 14 days, followed by a therapeutic dose of 2ml Nano-Selenium solution 0.25 or 0.5 mg/kg/day, respectively, via gavage tube once daily for 28 days. After forty-two days, the rats were ethically anesthetized by chloroform and euthanized. The right testes were excised, weighed, and

homogenized in buffer phosphate saline (1:10 parts, pH 7.4) using an Electrical Tissue Homogenizer (IKA, Germany); after that, the homogenate was centrifuged at 12,000 rpm at 4°C for 1 min and store at -20°C for antioxidant assays (19).

Animals monitoring

Measurement of oxidative stress parameters

In this study, the levels of three enzymes; Superoxide Dismutase (SOD), Glutathione peroxidase (GPX), and Malondialdehyde (MDA) were tested using ELISA Kits provided by SunLong Biotech Co., LTD to assay enzymes levels in the testicular tissues based on the interaction of pre-coated specific antibody on micrelisa strip plate wells with standards or samples.

Statistical analysis

The statistical package for social science (SPSS) (IBM SPSS Statistics for Windows, Version 23) was used for the analyses. One-way analysis of variance (ANOVA) was performed to summarize the results, and significant differences of P≤ 0.05 were applied.

Results

Characterization of *Eruca Sativa* extract

A. Total Flavonoid Content (TFC)

According to the experiment applied to estimate TFC, 0.9 g of the extract was found to be equivalent to 1 mg of Gallic acid.

B. Reducing Power Assay

A plot of log concentrations versus absorbance at 700 nm showed the right and downward shifting curve of *E. sativa* extract, compared to the vitamin C curve as a standard, which indicated that the extract's reducing activity is less than that of vitamin C within these selected concentrations, as shown in Figure (1).




Figure 1: Reducing power assay of *E. sativa* compared to vitamin C as a standard.

Characterization of Nano-Selenium

• Visual observation

Synthesis of SeNPs was indicated from the gradual conversion of the color of the extract from dark brown to light yellow and then reddish orange after about 12 hours of adding the acidic sodium selenite solution to the *Eruca Sativa* extract. This change can be ascribed to the aptitude of the extract to reduce selenium ions into SeNPs (Figure 2).




Figure 2:- Visual observation for Nano-selenium formation

A. UV-Vis Spectrum Analysis

Figure (3) shows the UV-Vis spectrum of SeNPs biosynthesized from *Eruca Sativa* extract. The spectrum of absorbance was measured from 220-1000 nm. A prominent absorption peak between 268 and 964 nm, with a maximum at 268 nm, is conclusive evidence of Nano selenium's production




Figure 3:- UV-visible spectrum of selenium nanoparticle.

B. Dynamic Light Scattering (DLS)

Particle size measurement

Table (1) shows that the average particle size of the synthesized Se-NPs varies from 39.4 to 124.6 nm. It also shows that the smallest particle size of the NPs is 39.4 nm, with PDI= 0.242, which was achieved using 1:2 of (Na₂SeO₃: *Eruca Sativa*) solution.

Zeta Potential

The produced Se-NPs had a zeta potential value of -56.57 mV. as shown in Figure (4), under optimal synthesis circumstances (1:2).

Table 1:- Poly disparity of nanoparticles and mean particle size

Sample	Sod.	Mean Particle size	Polydispersity
selenite: <i>E.sativa</i>			
1:2		39.4nm (minimum)	0.242
1:4		48.2nm	0.198
1:10		57.9nm	0.005
1:20		124.6nm (maximum)	0.304




Figure 4: Zeta potential of the synthesized Se-NPs made from sodium selenite with *Eruca Sativa* extract in the percentage of 1:2 (v/v).

C. Fourier transform infrared spectroscopy (FTIR)

To find out what biomolecules and plant extracts could interact with metal ions to create and stabilize selenium nanoparticles, FTIR spectroscopy was used. Figure 5A, B, and C shows the FTIR analysis results of *Eruca Sativa* extract. The spectral appearance of *Eruca Sativa* extract devoid of metal selenium is shown in (A). (B) is a representation of the sodium selenite spectrum, whereas the spectrum in (C) depicts the sample that includes selenium

metal in *Eruca Sativa* extract, hence displaying the peaks of both the control and test samples, as do the transmission peaks depicted in (B) and (A).




Figure 5: FTIR spectrum of (A) *Eruca Sativa* extract, (B) sod. selenite sol. and (C) selenium nanoparticles based- *Eruca Sativa* extract.

The plasma level of Malondialdehyde

The study showed a significant increase in the plasma level of MDA after exposing the animal to ketoconazole toxicity ($P<0.001$). MDA level was significantly higher in Gr B ($P<0.001$) compared to Gr A and this level stayed high even after treating the animal with *Eruca sativa* extract and sodium selenite (Gr BC and BD).

On the other hand, a prominent decline was noticed in Gr BE1 and Gr BE2. The levels were statistically insignificantly different compared to Gr A and stayed significantly lower than Gr B ($P<0.001$). Notably, the decline in MDA was prominently noticed after treating the animals with the Nanocomposite of selenium in the doses of 0.25 and 0.5mg/kg (BE1 and BE2), respectively. The MDA level of BE2 was significantly lower than that of groups B, BC, and BD ($P<0.001$) and was insignificantly different as compared to the negative control (Gr A) ($P<0.001$). At the same time, for BE1 there was a significant decrease compared to Gr B ($P<0.001$) but non-significant differences in comparison to Gr A, also, BC and BD.

The plasma levels of antioxidant enzymes

The study showed a prominent and statistically significant decline in the plasma levels of the antioxidant enzymes (Glutathione peroxidase and Superoxide dismutase) as measured in I.U/ml and pg/ml respectively, after having the animals exposed to ketoconazole toxicity ($P<0.01$) (Table 2). This decline persisted even after treating the ketoconazole-exposed rats with each of *Eruca sativa* extract and sodium selenite (Gr. BC and Gr.BD); the

plasma level stayed significantly lower than that of the negative control, with $P < 0.01$ and $P < 0.001$, respectively. Nevertheless, Gr. BD showed a statistically significant higher levels of plasma SOD and GPx as compared to Gr. B $P < 0.001$, and high significant decrease compared to Gr. A $P < 0.001$, with a significant increase $P < 0.01$, relative to Gr BC. Treatment with all forms of selenium nano

composite produced a statistically significantly higher level of these two enzymes than that of Gr. B, BC and BD, indicating the extract's potential to augment the body's antioxidant power ($P < 0.001$). At the same time, SOD and GPx in Gr BC and BD showed statistically lower levels than those in the negative control and higher levels than those in the positive control (Table 2).

Table 2: Levels of Malondialdehyde, superoxide dismutase and glutathione peroxidase in the homogenate of the testicular tissues of different treatment groups in rats

Group	MDA ng/g (tissue)	SOD I.U / g (tissue)	GPx pg/ g (tissue)
gr A	45.57 ± 3.01	7.80 ± 0.52	664.67 ± 23.05
gr B	71.38 ± 4.66 ***	4.42 ± 0.49 ***	338.67 ± 24.65 ***
gr BC	65.71 ± 5.17 ***	5.20 ± 0.45 ***	403.33 ± 37.43 ***
gr BD	63.43 ± 5.55 ***	5.83 ± 0.62 ***,**	497.33 ± 33.85 ***,**,##
gr BE1	55.48 ± 5.68 ***	7.52 ± 0.55 ***,###,☼☼	654.33 ± 29.21 ***,###,☼☼
gr BE2	43.33 ± 3.51 ***,###,☼☼	8.00 ± 0.44 ***,###,☼☼	662.67 ± 39.99 ***,###,☼☼

Gr. A & Gr B represent the negative and positive controls, respectively. Meanwhile, each of Gr BC and Gr. BD Represent the extract and pure sodium selenite, respectively. (Gr BE1 and BE2) were treated with selenium nanocomposite at 0.25 & 0.5 mg/kg, respectively. (*, ** & ***), (*, ** & ***), (#, ## & ###) and (☼, ☼☼ & ☼☼☼) signify a statistically significant difference as compared to groups A, B, BC and BD, with P values of $P < 0.05$, $P < 0.01$ and $P < 0.001$, respectively.

Discussion

In the reducing power assay test, Perlis Prussian stain's blue color appeared due to ferric ion reduction to ferrous ion in ferric ferrocyanide form. This test gives an idea about the potential of *Eruca Sativa* extract to donate electrons to free radicals (13).

According to the experiment that was performed to estimate total flavonoid content, 0.9 g of 1% *Eruca Sativa* extract is equivalent to 1 mg Gallic acid, which is compatible with Isra, 2022 results in which alcoholic extract and fresh rocket leaf contained high phenolic compounds and nearly equal amounts of total flavonoids content (0.81mg and 0.80 mg QE/g, respectively) (20).

The reddish-brown color is the most significant property of nanoparticle formation. Many researchers have previously obtained this result using different types of green plants (21) (Figure 2). Nanoparticle properties were characterized to confirm the formation of targeted Nano Selenium, using different techniques, including UV-Spectroscopy, Dynamic Light Scattering (DLS), and Fourier Transform Infrared Spectroscopy (FT-IR) (22).

The sample peak optical density, or absorbance, correlates linearly and directly to the concentration of nanoparticles in solution (23). According to Cittrarasu et al (2021) this SeNPs peak was created by the Surface Plasmon Resonance (SPR), which is induced when the electrons of metal nanoparticles interact with incoming photons (24), This result is compatible with many previous researches associated with various ranges of UV-visible maximum absorption peaks. Ramamurthy et al. found an absorbance peak ranging from 200–400 nm, with the maxima peak at 390 nm, when using

fenugreeek extract for SeNPs synthesis (25). Other SeNPs were synthesized by (26) using garlic cloves which reported an absorption peak at 260 nm. *Withania somnifera* was used to synthesize spherical SeNPs with a maximum absorbance of 310 nm, using Dynamic Light Scattering (DLS) (26). From the current study, it can be analyzed that there were uniformly dispersed particles, mostly in spherical Nano size, and no other peak was observed in the whole spectrum, meaning that Se^0 has been successfully formed (27).

DLS was performed to assess the particle size of different Nano solutions. The current study shows that, when increasing extract concentration in relation to sodium selenite; there was an increase in particle size diameter; the largest size 124.6 nm was related to the largest ratio of extract sample. This can be ascribed to the double quantity of extract solution available for reducing Na_2SeO_3 to Se^0 , resulting in more abrupt nucleation and faster growth of the nanoparticles. In some cases, small agglomerates were produced simultaneously (28). The least concentrated Nano sample has a much smaller mean size of approximately 39.4nm. However, it is characterized by a narrow size scatter compared to the sample with the largest extract ratio. The results were in line with the findings of Skandalis et al., in which Nanosilver was synthesized using Arbutus unedo extract, which gained a small particle size with the lowest concentrated extract sample.(29).

Polydispersity index (PDI) is a scaled and dimensionless index; when the value is higher than 0.7 it indicates that the sample has a very wide particle size distribution and that the DLS technique was probably not suitable to analyze it, while

monodisperse standard are mainly seen with the value smaller than 0.05 (30).

In general, zeta potential is the most important method to investigate the dispersion characteristics of nanoparticles and Nano fluids (31). It measures the potential differences between the charged ions on the nanoparticle surface that is opposite to the bulk fluid in which a particle is dispersed (32). The higher magnitude of potential differences around ± 60 mV exhibits higher electrostatic repulsion force and therefore increased stability, while agglomeration of particles may typically begin with a value lower than -15 mV. At the same time, zeta potential equals zero could represent the precipitation of the colloid into a solid (32). The Zeta potential value of the fabricated Se NPs was -56.57 mV, which indicated that the formed NPs were highly stable due to the negatively charged groups that surrounded them. The polydispersity index (PDI) value of the fabricated Se NPs was 0.242. This low value indicated that the formed Se NPs were mono-dispersed (33).

The functional groups present on the surface of the nanoparticles can be assessed by FTIR by measuring the chemical bond vibrational rates (34). Fig. (5) shows two absorption peaks located around 3248.13 & 3290.56 cm^{-1} that can be assigned as the absorption band of (O-H carboxylic acid), and one peak at 3371.57 cm^{-1} assigned as the absorption peak of (N-H) group. Also, FTIR spectrums depicted that IR bands in the regions 2935.66 & 2927.94 cm^{-1} are assigned for (C-H) asymmetric bending, in addition to three peaks at 1674.21, 1647.21 & 1604.77 cm^{-1} that may be due to (C=C) group (35). Similarly, two peaks at 1577.77 & 1573.91 cm^{-1} confirmed the presence of the (C=N) group. The absorption peaks at 1423.47, 1411.89 & 1400.32 cm^{-1} referred to the presence of the (O-H) group. Two peaks at 1149.57 & 1122.57 cm^{-1} referred to the (C-O-C) group, two peaks at 1083.99 & 1091.71 cm^{-1} to (C-N) group, and one peak at 829.35 cm^{-1} to (C-H) group (35).

In the present study, KET-induced testicular damage was associated with the reduction in both SOD and GPx peaks with the elevation in MDA level. These results are in harmony with previous studies that tested the ability of azoles to induce damage in different cellular models. In one study, Sertoli cells were exposed to different types of azoles and their combinations for 30min and 3h of incubation, azoles where elicited prooxidant properties associated with cell death, apoptosis, and necrosis (36). Also, marked changes in natural antioxidants (SOD and CAT) with oxidative damage to testicular lipids were associated with ketoconazole toxicity (36).

Although *Eruca Sativa* possesses powerful antioxidant properties as confirmed previously by total flavonoid content and reducing power assay, this effect was not powerful, as shown by the activities of the phyto-component conjugated selenium nanoparticles in reducing MDA and enhancing GPX and SOD level in the current study. This confirmed

the results of synergistic effects exerted by the bio-conjugation of Se-nanomaterial with plant extract in reducing their toxicities to biological systems and improving their therapeutic effectiveness (11)

The current results showed a significant decrease in MDA level in both BE1 and BE2 groups compared to Group B, in contrast to groups BC and BD which showed non-significant changes as compared to group B. These results confirmed the novel properties of the prepared nano solution-based *Eruca sativa* extract in reducing MDA, which can be considered as a diagnostic tool in male infertility (37). A previous study that tested the effects of nano selenium administration on ram sperm, previously exposed to the freeze/thawing process, showed a significantly increased quality of sperm by enhancing viability, motility, and membrane integrity, in addition to a significant reduction in the amount of MDA and damaged acrosome membrane (38).

The antioxidant enzymes, GPx, and SOD work together to scavenge the formed reactive oxygen species. Therefore, minor changes in normal constituents of these enzymes may disturb the body's defense systems with increasing the susceptibility to oxidative damage (39). Treatment with the Nanocomposite (Gr. BE1 and BE2) has elevated the level of GPX and SOD to be significantly higher than that obtained after treating the animals with sodium selenite or the extract alone (Gr. BD and BC) ($P < 0.001$); these results proved the novel characteristics of nanoparticles in crossing blood testes barrier (BTB) and maintaining the cellular redox, antioxidant capacity, and neutralizing ROS(40). The current results were compatible with other studies that have been previously published. Saber and Khalid, 2014, showed that Nano-selenium prepared by glutathione containing either 2 mg or 20 mg bovine serum albumin when administered in a dose of 0.5 mg/kg to male albino rats, would increase the defense mechanism and testicular antioxidant activity by enhancing the testicular GSH concentration, with reducing MDA concentration, with increasing catalase and SOD activity, as compared to controls and selenium groups (41). Furthermore, it has been reported that in addition to Se's role in forming selenoproteins, which help synthesize GPx enzymes, Se nanoparticles can up-regulate the expression of these enzymes through the formation of selenophosphate (41).

Conclusion

Nanotechnology is a revolutionary technology that has been gaining increased attention worldwide because of its potential role in medication delivery systems. Although selenium would be available for several metabolic, biological, and physiological functions, its nano forms prepared by *Eruca sativa* extract are unique in targeting the male reproductive system, by improving the antioxidant defense mechanism through its novel stability and small

particle size characteristics, suggesting a role against oxidative stress-related infertility induced by ketoconazole.

Authors' Declaration:

Conflicts of Interest: None

We hereby confirm that all the Figures and Tables in the manuscript are ours. Besides, the Figures and images, which are not ours, have been permitted republication and attached to the manuscript. -Authors sign on ethical consideration's approval-Ethical Clearance: The project was approved by the local ethical committee in (**College of Pharmacy /Al Nahrain University**) according to the code number (Nah. Co. Pha.14).

Author contributions:

Study conception: (Basman and Shayma'a) **Study design:** (Basman and Huda). Literature search: (**Study conception:** Basman and Shayma'a) Data acquisition: (Basman). Data analysis & interpretation: (Basman, Huda, and Shayma'a). Manuscript preparation: (Basman, Huda, and Shayma'a). Manuscript editing & review: (Basman, Huda, and Shayma'a).

References:

1. Alsaman ARS, Almashhedy LA, Hadwan MH. Effect of Oral Zinc Supplementation on the Thiol Oxido-Reductive Index and Thiol-Related Enzymes in Seminal Plasma and Spermatozoa of Iraqi Asthenospermic Patients. *Biol Trace Elem Res.* 2018; 184(2):340-9. <https://doi.org/10.1007/s12011-017-1215-8>
2. Landis-Piwowar KR, Iyer NR. Cancer chemoprevention: current state of the art. *Cancer Growth Metastasis.* 2014;7:19-25. <https://doi.org/10.4137/CGM.S11288>
3. Salman FS, Al-Qadhi HI, Al Kareem BA. N-acetyl cysteine's effect on semen parameters in a sample of Iraqi men with oligoasthenoteratozoospermia. *JFac Med Baghdad.* 2022;64(3):170-4. <https://doi.org/10.32007/jfacmedbagdad.6431938>
4. ASSUMAIDAE AAM, ALI NM, IBRAHEEM ZO, FADHIL AA, AL-HELLI KAA. Anti-Oxidoreductive stress Effects of Iraqi *Olea Europaea L.* Leaves Extract against Low Double Doses of Alloxan Induced Diabetes Mellitus in Rats. *SRP.* (2020),11(1): 292-302. [doi:10.5530/srp.2020.1.38](https://doi.org/10.5530/srp.2020.1.38)
5. Habbal H, Karabet F. Chitosan-(*Prunus avium*) Gum Nanocapsules Loaded with Orange Peel Extract: Nanocapsules Loaded with Orange Peel Extract. *Iraqi Journal of Pharmaceutical Sciences (P-ISSN 1683-3597 E-ISSN 2521-3512).* 2023; 32(1):194-201. <https://doi.org/10.31351/vol32iss1pp194-201>
6. Patra JK, Das G, Fraceto LF, Campos EVR, Rodriguez-Torres MDP, Acosta-Torres LS, et al. Nano based drug delivery systems: recent developments and future prospects. *J*

- Nanobiotechnology.* 2018;16(1):71. <https://doi.org/10.1186/s12951-018-0392-8>
7. Khamees EJ. Nanotechnology and Its Applications in Disease Diagnosis, Drug Delivery, and Medical Imaging. *Journal of the University of Babylon for Pure and Applied Sciences.* 2021:196-206. <https://www.journalofbabylon.com/index.php/JUBP/AS/article/view/3902>.
8. Hamza AA, Gamel M, Abdalla A, Abdalla Y, Amin A. *Gentiana lutea* attenuates hepatotoxicity induced by ketoconazole in rats by fortifying the cellular antioxidant defense system. *The Journal of Basic and Applied Zoology.* 2023;84(1):1-12. <https://doi.org/10.1186/s41936-022-00321-7>
9. Semet M, Paci M, Saïas-Magnan J, Metzler-Guillemain C, Boissier R, Lejeune H, et al. The impact of drugs on male fertility: a review. *Andrology.* 2017; 5(4):640-63. <https://doi.org/10.1111/andr.12366>
10. Al-Okaily BN. Role of alcoholic extract of *Rocket (Eruca sativa)* leaves on male reproduction of experimentally induced-oxidative stressed rats: Baraa Najim AL-Okaily and Ahmed Jasim Nowfel. *The Iraqi Journal of Veterinary Medicine.* 2015;39(2):47-54. <https://doi.org/10.30539/iraqijvm.v39i2.177>
11. Ibraheem ZO, Satar M, Abdullah NA, Rathore H, Tan YC, Uldin F, et al. Antioxidant and cardio protective effect of palm oil leaves extract (standardized ethanolic fraction) in rats' model of saturated fats induced metabolic disorders. *Pak J Pharm Sci.* 2014;27(1):1-9. [Pak J Pharm Sci 2014 27 1 1 9-libre.pdf \(dlwqxts1xzle7.cloudfront.net\)](https://doi.org/10.30539/iraqijvm.v39i2.177)
12. Ibraheem ZO, Sattar MA, Abdullah NA, Hassan R, Johns EJ. Toxicity, phytochemical content and antioxidant activity assessment studies for standardized ethanolic fraction of palm oil leaf extract. *Pharmacogn Commun.* 2012;2:21-30. <https://doi.org/10.5530/pc.2012.1.5>
13. Nhon Hoang TN, Phan TT, Lien Phan TK, Van Nguyen NH, Dao Dong TA, Anh Le TH. Phytochemical Screening, Extraction, and Determination of the Bioactivities of the Extract-Enriched Polyphenols and Saponins from *Musa balbisiana* Fruit. *Journal of Food Processing and Preservation.* 2023;2023. <https://doi.org/10.1155/2023/2581641>
14. Pyrzynska K, Sentkowska A. Biosynthesis of selenium nanoparticles using plant extracts. *Journal of Nanostructure in Chemistry.* 2021:1-14. <https://doi.org/10.1007/s40097-021-00435-4>
15. Naidu ECS, Olojede SO, Lawal SK, Rennie CO, Azu OO. Nanoparticle delivery system, highly active antiretroviral therapy, and testicular morphology: The role of stereology. *Pharmacol Res Perspect.* 2021;9(3):e00776. <https://doi.org/10.1002/prp2.776>
16. Anbu J, Mahendra SA, Sundar LM, Reddy RP,

- Nayak D. Evaluation of antifertility potential of Ketoconazole and assessment of its reversible effect on concomitant treatment with Metformin and Hydroxytyrosol in rats. *Asian Journal of Pharmacy and Pharmacology*. 2019;5(1):1-15. <https://doi.org/10.31024/ajpp.2019.5.1.1>
17. Abd-Elsalam RM, El Badawy SA, Ogaly HA, Ibrahim FM, Farag OM, Ahmed KA. *Eruca sativa* seed extract modulates oxidative stress and apoptosis and up-regulates the expression of Bcl-2 and Bax genes in acrylamide-induced testicular dysfunction in rats. *Environ Sci Pollut Res Int*. 2021;28(38):53249-66. <https://doi.org/10.1007/s11356-021-14532-y>
18. Khalaf AA, Ahmed W, Moselhy WA, Abdel-Halim BR, Ibrahim MA. Protective effects of selenium and nano-selenium on bisphenol-induced reproductive toxicity in male rats. *Hum Exp Toxicol*. 2019;38(4):398-408. <https://doi.org/10.1177/0960327118816134>
19. Vyas R, Kesari KK, Slama P, Roychoudhury S, Sisodia R. Differential Activity of Antioxidants in Testicular Tissues Following Administration of *Chlorophytum borivilianum* in Gamma-Irradiated Swiss Albino Mice. *Front Pharmacol*. 2021;12:774444. <https://doi.org/10.3389/fphar.2021.774444>
20. Dmour I. Physicochemical Characterization of Nanobiocomposites. *In Biocomposites-Recent Advances 2022 Dec 23. IntechOpen*. <https://doi.org/10.5772/intechopen.108818>
21. Jay V, Shafkat R. Synthesis of selenium nanoparticles using *Allium sativum* extract and analysis of their antimicrobial property against gram positive bacteria. *The Pharma Innovation*. 2018;7(9):262-6. <https://www.thepharmajournal.com/archives/2018/vol7issue9/PartE/7-9-34-545.pdf>
22. Titus D, James Jebaseelan Samuel E, Roopan SM. Chapter 12 - Nanoparticle characterization techniques. In: Shukla AK, Iravani S, editors. *Green Synthesis, Characterization and Applications of Nanoparticles*; Elsevier; 2019. p. 303-19. <https://doi.org/10.1016/B978-0-08-102579-6.00012-5>
23. Mohammed RA, Saleh GM, Mutlak FA. Investigate the Structural Properties of Silver Nanoparticles Produced by Bio Production of Green *Spinicia Oleracea* Leaf Extract and their Influence on Antibacterial Activity. *Iraqi Journal of Nanotechnology*. <https://www.iasj.net/iasj/download/Oee156c72c87b303>
24. Cittrarasu V, Kaliannan D, Dharman K, Maluventhen V, Easwaran M, Liu WC, et al. Green synthesis of selenium nanoparticles mediated from *Ceropegia bulbosa* Roxb extract and its cytotoxicity, antimicrobial, mosquitocidal and photocatalytic activities. *Sci Rep*. 2021;11(1):1032. <https://doi.org/10.1038/s41598-020-80327-9>
25. Ramamurthy C, Sampath KS, Arunkumar P, Kumar MS, Sujatha V, Premkumar K, et al. Green synthesis and characterization of selenium nanoparticles and its augmented cytotoxicity with doxorubicin on cancer cells. *Bioprocess Biosyst Eng*. 2013;36(8):1131-9. <https://doi.org/10.1007/s00449-012-0867-1>
26. Zambonino MC, Quizhpe EM, Jaramillo FE, Rahman A, Santiago Vispo N, Jeffryes C, et al. Green Synthesis of Selenium and Tellurium Nanoparticles: Current Trends, Biological Properties and Biomedical Applications. *Int J Mol Sci*. 2021;22(3). <https://doi.org/10.3390/ijms22030989>
27. Bibi I, Nazar N, Ata S, Sultan M, Ali A, Abbas A, et al. Green synthesis of iron oxide nanoparticles using pomegranate seeds extract and photocatalytic activity evaluation for the degradation of textile dye. *Journal of materials research and technology*. 2019;8(6):6115-24. <https://doi.org/10.1016/j.jmrt.2019.10.006>
28. Jasim AS, Salman ON. The Effect of Solvent Variation on Structural, Optical, and Electrical Properties of TiO₂ Films Prepared by Hydrothermal Method. *Journal of Applied Sciences and Nanotechnology*. 2023;3(2):59-69. <https://doi.org/10.53293/jasn.2023.5562.1191>
29. Skandalis N, Dimopoulou A, Georgopoulou A, Gallios N, Papadopoulos D, Tsipas D, et al. The Effect of Silver Nanoparticles Size, Produced Using Plant Extract from *Arbutus unedo*, on Their Antibacterial Efficacy. *Nanomaterials (Basel)*. 2017;7(7). <https://doi.org/10.3390/nano7070178>
30. Danaei M, Dehghankhold M, Ataei S, Hasanzadeh Davarani F, Javanmard R, Dokhani A, et al. Impact of Particle Size and Polydispersity Index on the Clinical Applications of Lipidic Nanocarrier Systems. *Pharmaceutics*. 2018;10(2). <https://doi.org/10.3390/pharmaceutics10020057>
31. Kmkm AM, Ghareeb MM. Natural Oil Nanoemulsion-Based Gel Vehicle for Enhancing Antifungal Effect of Topical Luliconazole. *J Fac Med Baghdad*. 2023; 65(1):65-73. <https://doi.org/10.32007/jfacmedbagdad.6512058>
32. Shim G, Jeong S, Oh JL, Kang Y. Lipid-based nanoparticles for photosensitive drug delivery systems. *J Pharm Investig*. 2022;52(2):151-60. <https://doi.org/10.1007/s40005-021-00553-9>
33. Sadeghi B, Gholamhoseinpoor F. A study on the stability and green synthesis of silver nanoparticles using *Ziziphora tenuior* (Zt) extract at room temperature. *Spectrochim Acta A Mol Biomol Spectrosc*. 2015;134: 310-5. <https://doi.org/10.1016/j.saa.2014.06.046>
34. Kadhim WK, Alshmmari MK, Al-Ethary MA. Synthesis and Characteristics of (PAV-PAVC-Ti) Nanocomposites. *Iraqi Journal of Nanotechnology, synthesis and application*. 2020; 1: 21-6.

<https://doi.org/10.47758/ijn.vi1.25>

35. Nandiyanto ABD, Oktiani R, Ragadhita R. How to read and interpret FTIR spectroscopy of organic material. Indonesian Journal of Science and Technology. 2019; 4(1):97-118.

<https://doi.org/10.17509/ijost.v4i1.15806>

36. Petricca S, Carnicelli V, Luzi C, Cinque B, Celenza G, Iorio R. Oxidative Stress, Cytotoxic and Inflammatory Effects of Azoles Combinatorial Mixtures in Sertoli TM4 Cells. Antioxidants. 2023; 12(6):1142.

<https://doi.org/10.3390/antiox12061142>

37. El-Sharawy ME, Hamouda M, Soliman AA, Amer AA, El-Zayat AM, Sewilam H, et al. Selenium nanoparticles are required for the optimum growth behavior, antioxidative capacity, and liver wellbeing of Striped catfish (Pangasianodon hypophthalmus). Saudi J Biol Sci. 2021; 28(12):7241-7.

<https://doi.org/10.1016/j.sjbs.2021.08.023>

38. Nateq S, Moghaddam G, Alijani S, Behnam M. The effects of different levels of Nano selenium on the quality of frozen-thawed sperm in ram. Journal of Applied Animal Research. 2020;48(1):434-9.

<https://doi.org/10.1080/09712119.2020.1816549>

39. Rajput VD, Harish, Singh RK, Verma KK, Sharma L, Quiroz-Figueroa FR, et al. Recent Developments in Enzymatic Antioxidant Defence Mechanism in Plants with Special Reference to Abiotic Stress. Biology (Basel). 2021;10(4).

<https://doi.org/10.3390/biology10040267>

40. Adhikari A, Mondal S, Das M, Ghosh R, Biswas P, Darbar S, et al. Redox Buffering Capacity of Nanomaterials as an Index of ROS-based Therapeutics and Toxicity: A Preclinical Animal Study. bioRxiv. 2021:2021.03.14.435286.

<https://doi.org/10.1101/2021.03.14.435286>

41. Asri-Rezaei S, Nourian A, Shalifar-Jalali A, Najafi G, Nazarizadeh A, Koohestani M, et al. Selenium supplementation in the form of selenium nanoparticles and selenite sodium improves mature male mice reproductive performances. Iran J Basic Med Sci. 2018; 21(6):577-85.

<https://doi.org/10.22038/IJBMS.2018.26023.6397>

How to Cite THIS Article

Antioxidant Effects of Selenium Nanoparticles Prepared from Eruca Sativa Extract on Ketoconazole –induced testicular Oxidative Damage in Male Rats. JFacMedBaghdad. 66 (1). Available from: <https://iqjmc.uobaghdad.edu.iq/index.php/19JFacMedBaghdad36/article/view/2174>

التأثيرات المضادة للأكسدة لجزيئات السيلينيوم النانوية المحضرة من مستخلص الجرجير على الكيتوكونازول – المسبب للأضرار التأكسدية في ذكور الجرذان

بسمان قاسم شريف¹, هدى ابراهيم الفاضلي¹, شيماء أحمد جمال²

¹ فرع الأدوية، كلية الطب، جامعة بغداد، بغداد، العراق

² فرع التشريح، كلية الطب، جامعة بغداد، بغداد، العراق

الخلاصة

خلفية البحث: أظهرت الدراسات الحديثة أن التأثيرات المضادة للخصوبة للكيتوكونازول يمكن التقليل منها عن طريق تناول مستخلصات نباتية تحتوي على جزيئات نانوية مضادة للأكسدة مع الدواء أو عن طريق تناوله بعد الدواء. إحدى الطرق الفعالة لتحسين خصائصه الطبية وتقليل سمية السيلينيوم هي صياغته على شكل جسيمات نانوية. ومع ذلك، لا تزال هذه استراتيجية صعبة.

هدف الدراسة: الهدف من هذه الدراسة هو تحليل تأثير السيلينيوم النانوي المحضر من مستخلص نبات الجرجير (Eruca Sativa) على معايير الإجهاد التأكسدي الخصوي في الجرذان المعالجة بالكيتوكونازول. ستركز الدراسة على تقييم إنزيم الجلوتاثيون بيروكسيداز، وفوق أكسيد ديسموتاز، والمالونديالدهيد كمؤشرات للإجهاد التأكسدي.

المواد والطرق: في هذه الدراسة، تمت إضافة محلول 1% وزن/حجم من المستخلص النباتي إلى محلول 10 ملي مولار سيلينيوم الصوديوم بنسب مختلفة، ووضع على محرك مغناطيسي في الظلام لمدة 12 ساعة عند درجة حرارة 50 درجة مئوية ودرجة الحموضة 9. تم بعد ذلك ترك المحلول لمدة 48 ساعة، وتم اختيار جزيئات السيلينيوم النانوية المثالية لمزيد من التوصيف. تم تقسيم ثمانية وأربعين فأراً إلى ست مجموعات تضم كل مجموعة ثمانية حيوانات. كانت المجموعة (أ) هي المجموعة الضابطة السلبية، في حين أعطيت المجموعة (ب) الكيتوكونازول عن طريق الفم بجرعة 50 ملغم/كغم لمدة أربعة عشر يوماً، يليها 200 ملغم/كغم من إيروكا ساتيفا، 0.5 ملغم/كغم من سيلينيوم الصوديوم عن طريق الفم، 25.0 ملغم/كغم من السيلينيوم النانوي عن طريق الفم و0.5 ملغم/كغم من السيلينيوم النانوي عن طريق الفم لمدة 28 يوماً على التوالي. وأخيراً، تم الموت الرحيم للحيوانات وتم تقييم المعلمات المضادة للأكسدة في الخصية.

النتائج: وقد لوحظت زيادات كبيرة في بيروكسيداز الجلوتاثيون وديسموتاز الفائق أكسيد (قيمة $p < 0.001$)، وانخفاض في مستويات المالونديالدهيد (قيمة $p < 0.001$) في المجموعات المعالجة بالجسيمات النانوية مقارنة بالمجموعة الضابطة.

الخلاصة: تظهر الأشكال النانوية المحضرة من مستخلص نبات الجرجير تأثيرات كبيرة مضادة للأكسدة على أنسجة الخصية، في حين أنها متاحة للعديد من الوظائف الأيضية والبيولوجية والفسولوجية.

مفتاح الكلمات: السيلينيوم، نانو سيلينيوم، سوبر اوكسايد دسموتاز، الجلوتاثيون بيروكسيداز، مالونديالدهيد