

Extraction and Identification of the Main Components of Cloves (*Syzygium aromaticum* L.) Oil Extract and its Antimicrobial Activity against Methicillin-resistant *Staphylococcus aureus* strain

DOI: <https://doi.org/10.32007/jfacmedbagdad.6622187>.Mays U. Hashim¹  , Orooba M.S. Ibrahim²  ¹Department of Physiology, Biochemistry and Pharmacology, College of Veterinary Medicine, Al-Qasim Green University, Babil, Iraq.²Department of Physiology, Biochemistry and Pharmacology, College of Veterinary Medicine, University of Baghdad, Baghdad, Iraq

©2024 The Author(s). Published by College of Medicine, University of Baghdad. This open-access article is distributed under the terms of the Creative Commons Attribution 4.0 International License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: Methicillin-resistant *Staphylococcus aureus* is widely recognized as a significant etiological agent responsible for infections around the world. One of the biggest problems in world health care is antibiotic resistance to the MRSA strain. The use of herbal medicines is one of the promising techniques for countering bacterial resistance to antibiotics.

Objectives: The study is designed to investigate the chemical composition of clove oil extract and its *in-vitro* antibacterial activities against MRSA.

Methods: The clove oil extract was obtained by using hydro-distillation by Clevenger apparatus. After that, phytochemical analysis was done to determine the secondary metabolites by Chromatography-Mass Spectrometry. *In-vitro* antimicrobial activity of clove oil extract against Methicillin-resistant *Staphylococcus aureus* was carried out by agar well diffusion method, the broth microdilution method, and *in-vitro* time-kill curve kinetic. Least significant difference –LSD test (Analysis of Variation-ANOVA) was used to significant compare between means of results in this study.

Results: The results of this study revealed that the extraction percentage of the clove yielded 50%. The Chromatography-Mass Spectrometry results of the clove oil extract analysis showed that caryophyllen at 28.9%, Humulene at 21.6% and eugenol at 13.06% were the primary bioactive ingredients of the prepared extract. Furthermore, the minimum inhibitory concentration (MIC) and the minimum bactericidal concentration (MBC) of clove oil extract against *Methicillin-resistant Staphylococcus aureus* were found to be 2.5µg/ml and 5.0 µg/ml respectively. Time killing curve of 2xMICs and 4xMICs of clove extract achieved the highest significant bactericidal effect (P≤0.05) in comparison to other concentrations.

Conclusions: The clove oil extract exhibited good *in-vitro* antibacterial properties and this can be attributed to the presence of phenolic compounds such as caryophyllene, humulene and eugenol.

Keywords: Antimicrobial, Caryophyllen, Clove extract, *Staphylococcus aureus* (MRSA), Time killing curve.

J Fac Med Baghdad
2024; Vol.66, No. 2
Received: Jul., 2023
Revised: Sept. 2023
Accepted: Jan 2024
Published: Jul. 2024

Introduction

Multidrug-resistant bacteria causing infectious diseases were responsible for significant mortality, particularly in developing countries (1). High prevalence of methicillin-resistant *Staphylococcus aureus* strains (MRSA) was recorded in both health-care and community environments (2,3,4). To develop new drugs to treat multidrug-resistant pathogens, the search for novel antibacterial compounds, including herbal products, has increased quickly (5,6). Traditional treatments frequently use pharmaceuticals made from aromatic herbs to treat bacterial infections (7). Numerous investigations claim that utilizing essential oils can help to decrease antibiotic-resistant bacteria (8).

These oils have a wide range of biological and pharmacological effects and are very volatile, lipophilic, and hydrophobic (9). *Syzygium aromaticum* (family- *Myrtaceae*), more popularly known as clove, is indigenous to the Indonesian islands. Almost there is thirty compounds have been identified in Clove Essential Oil Composition (CEO), eugenol is the main component, composing at least half of its components. Caryophyllene, Eugenyl acetate, and humulene compose the remainder, which ranges from 10%–40%. The remainder is less than 10% which is considered minor elements(10). It is now grown all over the world as a flavoring ingredient, a medicine, and an ingredient in perfumes (11,12). Due to its antibacterial abilities, it is frequently used as a food preservative (13). Analgesic, antioxidant, anti-inflammatory,

* Corresponding author's Email:
maysudayhashim@vet.uoqasim.edu.iq

anesthetic, and insecticidal action has also been documented in addition to its antibacterial property (14) Additionally, the antibacterial activity of clove oil have been studied against a large number of multi-resistant *Staphylococcus* and oral pathogens(15). The development of innovative antimicrobial treatment agents is necessary due to the high prevalence of bacterial strains that are resistant to many drugs. Thus, the goal of the current study was to determine the chemical composition as well as the *in-vitro* antibacterial effect of clove oil extracts against pathogenic MRSA bacterial strains.

Materials and Methods

Collection of plant

The origin of clove (*S.aromaticum*) utilized in this study is from a local market in Baghdad, Iraq, and classified by the National Center For Herbal Medicine and Al-Razi Center for Medical Herbs. Clove buds were cleaned with 5% sodium hypochlorite solution (NaOCl), rinsed three times with distilled water, and then kept to dry. Using a mechanical mortar, the dried plant was ground into powder.

Extraction of clove oil (*Syzygium aromaticum*)

Clove oil was extracted from dry buds using hydro-distillation by Clevenger apparatus. After drying with sunlight, it was placed in a 1000 ml glass flask then 500 ml of distilled water was added. The flask was connected to the apparatus and operated for three hours. The flask's temperature was first raised to about 80°C and then gradually increased to 100°C. The oil was collected after isolating the water, and to increase the water disposal a little of anhydrous sodium sulfate was used (16).

Extraction percentage yield of essential Clove oil

The collected cloves sample was used to extract the essential oil by hydrodistillation method. The extraction yield has been accomplished by the equation below (17):

$$\text{Oil yield \%} = \frac{\text{Volume of oil extracted}}{\text{weight of the sample}} * 100$$

GC-MS analysis of *S. aromaticum* Oil extract

To identify the active components of clove extraction, Gas chromatography-mass spectrometry (GC-MS) was used to study the phytochemicals of the clove (*S. aromaticum*) extracts. By injecting 1 L of the sample (0.1% in absolute methanol) and operating in scan mode on the GC/MS Thermo Trace GC Ultra / TSQ Quantum GC-MS, the GC/MS analysis was carried out. Using an Agilent HP- 5ms Ultra Inert capillary column (30 m 0.25 m film thickness), the phytochemical investigation was conducted. The rates of the four ramps were as follows: ramp 1 was 60 °C hold to 3 min, ramp 2 was 60 °C to 180 °C hold for 7 min, ramp 3 was 180 °C to 280 °C holding to 8 min, and Ramp 4 was 280 °C holding to 3 min. The following describes the operation's conditions: The carrier gas was helium, with a 99.99% purity, and the injector and detector were 250 °C hot. Comparing the

results of the GC-MS analysis with the reference retention time and spectral mass data from the NIST database allowed the chemical components of the clove bud extract to be identified.

Bacteriological Examination

Test Organism

The Methicillin resistance *staphylococcus aureus* (MRSA) was obtained from the Physiology, Biochemistry, and Pharmacology Department/ College of Veterinary Medicine /University of Baghdad.

Activation and Maintenance of MRSA

Ten mL of brain heart infusion agar slants were used to activate bacterial cultures in screw-capped tubes and then placed in the incubator at 37 °C for 24 hrs. For maintenance of isolates, the bacterium was cultured on brain heart infusion agar and kept at 4°C, then the bacterium was activated every 14 days. These bacteria were established for microscopic morphological, cultural, and biochemical studies (18).

Preparation of standard bacterial suspension

By comparing to the Standard McFarland solution (0.5), the quantity of MRSA bacteria in each milliliter of the stock suspension was standardized (18). Briefly, bacterial suspension equivalent to 0.5 McFarland (1.5×10^8 CFU /ml) was arranged from overnight bacterial culture. The absorbance of this index was 0.136 as noted by the spectrophotometer at a wavelength of 450 nanometers. About 0.1 ml of the prepared bacterial suspension was diluted in 14.9 ml of Mueller-Hinton broth and incubated at 37 °C for 1 hr. to obtain 10^6 CFU/ml bacterial suspensions to prepare bacterial suspension for the time-kill curve kinetic assay in order to bring MRSA bacteria to the logarithmic phase of bacterial growth (19).

Measurement of Antimicrobial Activity of clove oil extract:

Agar well diffusion method:

The agar well diffusion method was carried out to evaluate the antibacterial activity of *S.aromaticum* extract against MRSA according to (20). Standardized bacterial suspension (1.5×10^8 cfu/ml) of *S. aureus* was carefully mixed with sterile Mueller Hinton agar. Twenty-five ml of this agar was dispersed into sterile Petri dishes and left for 10 minutes at room temperature to dry, and six mm. diameter wells were bored in the agar. The *S.aromaticum* extract was reconstituted in distilled water to a concentration from $5\mu\text{g/ml}$ to $0.312\mu\text{g/ml}$ and then $100\mu\text{l}$ was added to wells. The plates were incubated at 37 C° for 24 hours after allowing the extract to diffuse into the agar at room temperature. Three plates were made for each concentration with negative control using buffer phosphate and the strength of the clove oil extract was dictated by measuring the inhibition zone diameter around every well against the tested bacteria and used as positive bioactivity compared with negative control. Standard error and mean were calculated.

Microtiter plate Dilution

Clove oil extract (10 mg/ml) was made in Mueller-Hinton broth, from this broth, two folds dilution was downgraded from 40 µg for clove oil in U- shape (200 µl well capacity) 96 well micro-titer plate. Each well was inoculated with 100 µl of 1.5×10^8 CFU/ml *S. aureus* and incubated at 37°C for 24hrs. For colorimetric identification of bacterial growth, 15µl of 0.125% triphenyl tetrazolium chloride dye (TTC) which was used as an indicator of cellular viability was added to each well of the test and re-incubated for two hours to determine MIC by observing whether or not the red color that results from the reductions of TTC (colorless) to formazan (red) develops (21). While the minimum bactericidal concentration (MBC) was determined by subculturing 50 µl from the well that showed no apparent growth (clear) onto fresh nutrient agar plates. After the incubation period, if there was no growth this concentration was taken as MBC which is considered the lowest concentration of extracts that kill the bacteria (22).

Time Kill Curve Kinetic Assay

The time-kill curve assay of clove extract against MRSA bacteria was done according to the procedure described by (19). Briefly, bacterial suspension was prepared as mentioned before to obtain 10^6 CFU/ml bacterial suspensions, and the Clove extract had been dissolved in Mueller-Hinton broth to prepare 10 mg/ml stock solution. After that, clove extract concentrations from 4x MIC to 0.25x MIC were prepared. Bacterial colonies were calculated at 0, 1, 2, 4, 6, and 24 hr. through the incubation time by making serial dilutions and spreading of 20 µl of each dilution on Mueller-Hinton agar plate (triplicate); colonies range 30-300 CFU/plate was accepted (19).

The trapezoidal method was used to estimate the area under the time-kill curve of the concentration of clove extract as below (23):

$$AUC_{kill} = \sum \left(\frac{\log(C_n) + \log(C_{n+1})}{2} \cdot \Delta t \right)$$

Where the AUC_{kill} is the area under the killing curve of clove extract, $\log(C_n)$ is the logarithm of a specific concentration at a specific time, $\log(C_{n+1})$ is the logarithm of the next concentration while (Δt) resembles the difference between their times.

Statistical Analysis

The Statistical Analysis System- SAS program was used to detect the effect of different factors on study parameters. Least significant difference –LSD test (Analysis of Variation-ANOVA) was used to significantly compare between means in this study.

The authentication of the plant

The clove that utilized in this study was classified by the National Center For Herbal Medicine and Al-Razi Center for Medical Herbs as *Syzygium aromaticum* L. family of *Myrtaceae* to be used as oil extract in this study.

Extraction of essential Clove oil

The collected cloves sample was used to extract the essential oil by hydrodistillation method. The oil

obtained was yellow in color as show in (Figure 1) with the highest extraction yield of 50%.




Figure1: Clove oil extract

Results:

Gas chromatography-mass spectrometry (GC-MS)

Seventeen peaks related to separate components were obtained from gas chromatography–mass spectrometry (GC/MS) in (*S. aromaticum*) extract by hydrodistillation method (Figure 2).




Figure 2: Gas chromatogram of clove oil extract

The main constituents were caryophyllene with 28.9%, eugenol 13.06%, and Humulene 21.6% from the GC-MS database. Additionally, other compounds were listed in (Table 1).

Table 1: Phytochemical analysis of *S. aromaticum* essential oil.

Compounds	Retention time	Percentage of total %
Caryophyllene	13.9	28.9
Humulene	14.5	21.6
Eugenol	12.7	13.06

Antibacterial activity of clove oil extract Agar well diffusion method

Different concentrations of clove oil extract were used in agar well diffusion methods, resulting in different sizes of inhibition zones against methicillin resistance *Staphylococcus aureus*. The sizes of inhibition zones were different according to the concentration of the Clove oil extract. Results indicated that MRSA bacteria were sensitive significantly ($P < 0.05$) to clove oil extract in a concentration-dependent manner 5, 2.5, 1.25, 0.625, and 0.312 µg/ml. Increasing the diameter measurement of the zone of inhibition in MRSA

growth was proportionally related to clove oil extract concentrations, (Table 2 and Figure 3).

Table 2: Antibacterial activity of Clove oil extract in different concentrations against MRSA (measured as the diameter of the inhibitory zone in millimeters).

Group	Con.(µg/ml)				
S	5µg/ml	2.5µg/ml	1.25µg/ml	0.625µg/ml	0.312µg/ml
Conc.					
Clove oil extract	18.0±0.75 A a	15.0±0.58 B a	13.0±0.61 BC a	11.0±0.52 C a	9.0±0.38 C a
Buffer phosphate	0.0±0.0 A b	0.0±0.0 A b	0.0±0.0 A b	0.0±0.0 A b	0.0±0.0 A b
LSD value	3.337*				

- Values represent mean ± S.E
- Different capital letters mean significant ($P < 0.05$) results between different concentrations.
- Different small letters mean significant ($P < 0.05$) results between buffer and clove oil extract.
- *LSD: least significant difference




Figure 3: Susceptibility of MRSA strain to different concentrations of clove extract

Microtiter plate Dilution

The Minimum Inhibitory Concentration plays a key role in the determination of an antibacterial potency (24). The microdilution assay was used to determine MIC. The results showed that the concentration of 2.5 µg/ml of clove oil was effective in preventing MRSA from growing, this concentration was shown to have a positive value that inhibited the growth and 5.0 µg/ml of clove oil extract killed MRSA so they were considered as the (MBC). MBC value was tested in micro-dilution assay by sub-culturing 50 µl from the well that showed no apparent growth (clear), if there was no growth this concentration was taken as MBC.

Time Kill Curve Kinetic Assay

The time-kill curve kinetics of clove oil extract is based on the highest MIC recorded from the micro-dilution assay which was 2.5µg/ml for the MRSA strain. The concentrations used *in vitro* study were 0.25x MIC, 0.5x MIC, 1x MIC, 2x MICs, and 4x MICs.

The clove oil extract at a concentration of (4xMIC, 2xMIC) showed significant killing activities both at 6 and 24 hrs. respectively by reducing ≥ 3 log₁₀ of the total number of CFU/ml in comparison to other concentrations used as shown in (Figure 4).




Figure 4: Time kill curve kinetics of clove extract against MRSA

The area under the time of killing curve of clove extract was calculated and compared to the control inoculum growth rate, and the difference in the area under the curve values among different treatments was set as an endpoint whereas the lowest area under the curve refers to the highest bactericidal effect as reported in the (Table 3). The results showed that both 2xMICs and 4xMICs achieved the highest significant bactericidal effect ($P \leq 0.05$) in comparison to other treatments. The 1xMIC concentration achieved a purely bacteriostatic effect ($P \leq 0.05$) in comparison to all concentrations and control groups; both 0.5xMIC and 0.25xMIC failed to achieve a significant bacteriostatic or bactericidal effect ($P \geq 0.05$) in comparison to 1xMIC, 2xMIC and 4xMIC.

Table 3: Area under the time-kill curve of clove extract against MRSA (h*log₁₀ CFU/ml).

Antibacterial	Control	0.25xMIC	0.5xMIC	1xMIC	2xMIC	4xMIC
Clove extract	125.3 ±0.31 A	111.5 ±0.3 A	86.81 ±0.30 B	57.22 ±0.19 C	29.8 ±0.5 D	17.2 ±0.5 D
MIC=2.5µg/ml	13.3*					
SD						

- Values represent mean ± S.E
- Different capital letters denoted a significant difference ($p \leq 0.05$) among the groups.
- * LSD: least significant difference

Discussion

Clove oil extraction using hydro-distillation (water distillation) by Clevenger apparatus gave a bright yellow color oil and brown color extract with a typical clove oil smell. The extraction yield of clove extract was 50% as the amount of clove buds that were used in the current procedure was 200 g. However this outcome is nearly in line with the outcomes of the study of Ishaq and his colleagues(2019) who discovered that when clove bud powder was extracted using a soxhlet equipment, the yield of hexane extract was 48.84% (25). Importantly, one of the reason of

the resulting higher percentage of extraction may belong to the reduction in the particle size as reported by Ratri and his colleagues (2020) as they suggested the percentage yield of clove extract is influenced by the cloves' particle size, with smaller clove particles producing higher extraction yields (26). However, the extraction procedure can also be influenced by several other factors, such as the quality and freshness of the clove buds, the temperature and pressure used during the distillation process, and the duration of the distillation process (27, 28).

To identify the active components of clove extraction, GC/MS was used and the main compounds of the *S. aromaticum* extract were caryophyllen, humulene, and eugenol. According to reports, these substances make up the majority of the clove buds' active ingredients which have a therapeutic use such as using as analgesic, antioxidant, anti-inflammatory, anesthetic, and antibacterial agents (29). In this study ground clove was used and found β -caryophyllene be the most abundant since the volatile profile from whole buds showed a different pattern when compared to the volatile composition of ground clove. And this is in agreement with Gaspar and his colleagues (2018) who approved that the most prevalent compound is β -caryophyllene. with (49.31% concentration) as they compared between compositions from whole and ground clove (30). However, other studies, have found a difference in the majority of compounds, such as the study of Lee (2009) who has shown that eugenol, β -caryophyllene, 2-propanone, and methylhydrazone are the main composites of (*S. aromaticum*) essential oil (31).

The anti-MRSA potential of clove oil extract has been reported and has highlighted the requirement to investigate bioactive components (32). The clove extract exerted the highest antimicrobial efficacy against the pathogenic bacterial strains (MRSA), which may be assigned to the high percentage of a phenolic compound of clove oil extract as presented in GC-MS results and this complies with the result of Alanazi and his colleagues (2022) who demonstrated the lowest concentration of clove oil that inhibit the growth of MRSA found to be 2.5 μ L/ML (33).

The phytochemical constituents that are present in clove oil extract are responsible for the antibacterial activity. Eugenol is one of the bioactive components that may have considerably contributed to the antibacterial effects of the clove oil (34). Humulene is another component that present in clove oil similar to caryophyllene with antibacterial properties (35). However, the minor components could potentially contribute by combining them with other main components to limit MRSA development. The oil's hydrophobic properties may also aid in interactions with the outer cytoplasmic membrane of MRSA, which impairs the integrity and functionality of the cell membrane (36). Time killing-curve kinetic is a combined and extensive tool to assess both bacteriostatic and bactericidal effects of the

antibiotics; it depends on the change in the logarithmic number of bacterial colonies through the defined chronological pattern (19). More accurate descriptions of antimicrobial activity are provided by a measure of bacterial killing (kill kinetics) than by the MIC, and it has also shown better sensitivity developments to physicians than disc diffusion methods (37, 38)

The current time-kill kinetic outcomes required a little more time to completely eradicate the bacteria when compared with the study conducted by Mandal and his colleagues (2011) who found that clove extract showed significant killing activities against MRSA, both at 3 and 6 hrs at a concentration of 256 μ g/ml ($1 \times$ MIC) (38). These differences are possibly caused by a variety of bacterial species used, the concentration of the antibacterial used, and the method used.

Conclusion: The clove oil extract at different concentrations had antimicrobial activity against MRSA bacteria. The GC-MS analysis of the clove oil showed the presence of 17 volatile components that have been previously reported to possess antibacterial effects. Hence the antibacterial properties demonstrated by the clove oil extract can be attributed to the compounds identified caryophyllen, humulene, and eugenol. Therefore, the clove oils may be used in the medicinal formulation of antimicrobial drugs.

Authors' declaration:

Conflicts of Interest: The authors declare no conflict of interest. We confirm that all the Figures and Tables in the manuscript are mine/ ours. Besides, the Figures and images, which are not mine /ours, have been given permission for re-publication and attached to the manuscript. ethical consideration's approval-Ethical Clearance: The project was approved by the local ethical committee in (collage of Veterinary Medicine/ University of Baghdad) according to the code number (1604 in 26-7-2023).

Author contributions:

Study conception & design: (Orooba MS). Literature search: (Mays Uday & Orooba MS). Data acquisition: (Mays Uday). Data analysis & interpretation: (Mays Uday & Orooba MS). Manuscript preparation: (Mays Uday & Orooba MS). Manuscript editing & review: (Mays Uday) .

References

- 1- Taati M, Khoshbayan A, Chegini Z, Farahani I, Shariati A. Bacteriophages, a new therapeutic solution for inhibiting multidrug-resistant bacteria causing wound infection: lesson from animal models and clinical trials. *Drug design, development and therapy*. 2020 May 15;1867-83. DOI: [10.2147/DDDT.S251171](https://doi.org/10.2147/DDDT.S251171)
- 2- Abdelraheem WM, Khairy RM, Zaki AI, Zaki SH. Effect of ZnO nanoparticles on methicillin, vancomycin, linezolid resistance and biofilm

- formation in *Staphylococcus aureus* isolates. *Annals of Clinical Microbiology and Antimicrobials*. 2021 Aug 21;20(1):54. <https://doi.org/10.1186/s12941-021-00459-2>.
- 3- Kanaan MH, AL-Shammary AH. Detection of methicillin or multidrug resistant *Staphylococcus aureus* (MRSA) in locally produced raw milk and soft cheese in Baghdad markets. *The Iraqi Journal of Veterinary Medicine*. 2013 Dec 31;37(2):226-31. <https://doi.org/10.30539/ijvm.v37i2.1382>
- 4- Yassin MT, Al-Askar AA, Mostafa AA, El-Sheikh MA. Bioactivity of *Syzygium aromaticum* (L.) Merr. and LM Perry extracts as potential antimicrobial and anticancer agents. *Journal of King Saud University-Science*. 2020 Dec 1;32(8):3273-8. <https://doi.org/10.1016/j.jksus.2020.09.009>
- 5- Sannathimmappa MB, Nambiar V, Aravindakshan R. Antibiotics at the crossroads—Do we have any therapeutic alternatives to control the emergence and spread of antimicrobial resistance?. *Journal of Education and Health Promotion*. 2021;10. <https://doi.org/10.4103%2Fjehp.jehp.557.21>
- 6- Abdul-Jalil TZ. Evaluation of the active constituents, Antioxidant and Antimicrobial Activities of Iraqi *Euonymus japonicus* leaves using Ethyl Acetate Extract: Polyphenolic Characterization, Antioxidant and Antibacterial Activity of Iraqi *Euonymus japonicus* L. Ethyl Acetate Extract. *Journal of the Faculty of Medicine Baghdad*. 2023 Jul 1;65(2). <https://doi.org/10.32007/jfacmedbagdad.1991>
- 7- Yazarlu O, Iranshahi M, Kashani HR, Reshadat S, Habtemariam S, Iranshahi M, Hasanpour M. Perspective on the application of medicinal plants and natural products in wound healing: A mechanistic review. *Pharmacological research*. 2021 Dec 1; 174:105841. <https://doi.org/10.1016/j.phrs.2021.105841>
- 8- Mohd NA, Mohd MI, Manickam S, Sun X, Goh BH, Tang SY, Ismail N, Abdull Razis AF, Ch'ng SE, Chan KW. Essential oils and plant extracts for tropical fruits protection: From farm to table. *Frontiers in Plant Science*. 2022 Sep 29;13:999270. <https://doi.org/10.3389/fpls.2022.999270>
- 9- Oliveira TA, Santiago MB, Santos VH, Silva EO, Martins CH, Crotti AE. Antibacterial activity of essential oils against oral pathogens. *Chemistry & Biodiversity*. 2022 Apr; 19(4):e202200097. <https://doi.org/10.1002/cbdv.202200097>
- 10- Haro-González JN, Castillo-Herrera GA, Martínez-Velázquez M, Espinosa-Andrews H. Clove essential oil (*Syzygium aromaticum* L. Myrtaceae): Extraction, chemical composition, food applications, and essential bioactivity for human health. *Molecules*. 2021 Oct 22;26(21):6387 <https://doi.org/10.3390/molecules26216387>
- 11- Ali BM, Ibrahim O. Antifungal Activity of Clove (*Syzygium aromaticum*) Essential Oil Extract against Induced Topical Skin Infection by *Candida albicans* in Mice In-vivo. *The Egyptian Journal of Hospital Medicine*. 2023 Apr 1;91(1):3855-61. <https://dx.doi.org/10.21608/ejhm.2023.293468>
- 12- Batiha GE, Beshbishy AM, Tayebwa DS, Shaheen HM, Yokoyama N, Igarashi I. Inhibitory effects of *Syzygium aromaticum* and *Camellia sinensis* methanolic extracts on the growth of *Babesia* and *Theileria* parasites. *Ticks and tick-borne diseases*. 2019 Aug 1;10(5):949-58. <https://doi.org/10.1016/j.ttbdis.2019.04.016>
- 13- Kumar P V, Shams R, Singh R, Dar AH, Pandiselvam R, Rusu AV, Trif M. A comprehensive review on clove (*Caryophyllus aromaticus* L.) essential oil and its significance in the formulation of edible coatings for potential food applications. *Frontiers in Nutrition*. 2022 Sep 15; 9:987674. <https://doi.org/10.3389/fnut.2022.987674>
- 14- Charoonratana T. Chapter 5 - Clove (*Syzygium aromaticum*). *Clove (Syzygium Aromaticum)*. 2022 Jan 1:49-65. <https://doi.org/10.1016/B978-0-323-85177-0.00001-X>
- 15- Mishra RP, Kalyani S. Antimicrobial activities of *Syzygium aromaticum* L.(Clove). *Int Res JBiological Sci*. 2014;3(8):22-5. <https://www.academia.edu/download/89149159/4.IS-CA-IRJBS-2014-42.pdf>
- 16- Taraj K, Andoni A, Fatos YL, Ariana YL, Hoxha R, Llupa J, Malollari I. Spectroscopic investigation of *Syzygium aromaticum* L. oil by water distillation extraction. *Journal of International Environmental Application and Science*. 2020; 15(2):122-6. https://dergipark.org.tr/en/pub/jieas/issue/54918/705980#article_cite
- 17- Tambe E, Gotmare S. Qualitative estimation of chemical composition of five different clove oils (*Syzygium aromaticum*) by GCMS. *International Journal of Multidisciplinary Educational Research*. 2020;9(6):5. https://www.researchgate.net/publication/344197071_qualitative_estimation_of_chemical_composition_of_five_different_clove_oils_syzygium_aromaticum_by_gcms
- 18- Ibrahim OM, Saliem AH, Salih SI. Antibacterial activity of silver nanoparticles synthesized by Cinnamon zeylanicum bark extract against *Staphylococcus aureus*. *Al-Anbar Journal of Veterinary Sciences*. 2016;9(1):22-36. <https://www.iasj.net/iasj/article/113248>
- 19- Shwaish MM, Ibrahim O. Pharmacodynamics analysis of meropenem against *klebsiella pneumonia* isolated from covid-19 patient. *Biochemical & Cellular Archives*. 2022 Apr 1;22(1). <https://Pharmacodynamics+analysis+of+meropenem+against+klebsiella+pneumonia+isolated+from+covid-19+patient&bmG>
- 20- Salih AN, Ibrahim OM, Eesa MJ. Antibacterial activity of biosynthesized silver nanoparticles against *Pseudomonas aeruginosa* In-vitro. *The Iraqi Journal of Veterinary Medicine*.

2017;41(1):60-5 .

<https://doi.org/10.30539/iraqijym.v41i1.81>

21- Mohammed SA, Ibrahim O. Pharmacodynamics analysis of fosfomycin against multidrug resistant *e. coli* o157: h7 isolated from urinary tract infection. *Biochemical & Cellular Archives*. 2022 Apr 1; 22(1).

https://scholar.google.com/scholar?cluster=7016470094531529743&hl=ar&as_sdt=0,5

22- Al-Mizraqchi AS. Anti-microbial activity of Green Tea Extracts and Nicotine on the Growth, Biofilm Formation of *Sali-vary Mutans Streptococci* (In-vitro study). *Journal of the Faculty of Medicine Baghdad*. 2023 Jul 1; 65(2).

<https://doi.org/10.32007/jfacmedbagdad.2024>

23- Chiou WL. Critical evaluation of the potential error in pharmacokinetic studies of using the linear trapezoidal rule method for the calculation of the area under the plasma level-time curve. *Journal of pharmacokinetics and biopharmaceutics*. 1978 Dec; 6:539-46 .

<https://doi.org/10.1007/bf01062108>

24- Jasim MA, Saeed OM. pharmacodynamic properties of meropenem against *staphylococcus aureus*. *Biochemical & Cellular Archives*. 2021 Apr 1;21(1).

https://www.researchgate.net/publication/351230283_PHARMACODYNAMIC_PROPERTIES_OF_MEROPENEM_AGAINST_STAPHYLOCOCCUS_AUREUS

25- Ishaq A, Syed QA, Khan MI, Zia MA. Characterizing and optimizing antioxidant and antimicrobial properties of clove extracts against food-borne pathogenic bacteria. *International Food Research Journal*. 2019 Jul 1;26(4):1165-72 .

[http://www.ifrj.upm.edu.my/26%20\(04\)%202019/6%20-%20IFRJ181176.R1-Final.pdf](http://www.ifrj.upm.edu.my/26%20(04)%202019/6%20-%20IFRJ181176.R1-Final.pdf)

26- Ratri PJ, Ayurini M, Khumaini K, Rohbiya A. Clove oil extraction by steam distillation and utilization of clove buds waste as potential candidate for eco-friendly packaging. *Jurnal Bahan Alam Terbarukan*. 2020 20; 9(1):47-54 .

<https://doi.org/10.15294/jbat.v9i1.24935>

27- Selles SM, Kouidri M, Belhamiti BT, Ait Amrane A. Chemical composition, in-vitro antibacterial and antioxidant activities of *Syzygium aromaticum* essential oil. *Journal of Food Measurement and Characterization*. 2020 ;14(4):2352-8 <https://doi.org/10.1007/s11694-020-00482-5>

28- Gonzalez-Rivera J, Duce C, Campanella B, Bernazzani L, Ferrari C, Tanzini E, Onor M, Longo I, Ruiz JC, Tinè MR, Bramanti E. In-situ microwave assisted extraction of clove buds to isolate essential oil, polyphenols, and lignocellulosic compounds. *Industrial Crops and Products*. 2021 = 1; 161:113203 <https://doi.org/10.1016/j.indcrop.2020.113203>

29- Kumar PV, Shams R, Singh R, Dar AH, Pandiselvam R, Rusu AV, Trif M. A comprehensive review on clove (*Caryophyllus aromaticus* L.)

essential oil and its significance in the formulation of edible coatings for potential food applications. *Frontiers in Nutrition*. 2022 = 15; 9:987674 .

<https://doi.org/10.3389/fnut.2022.987674>

30- Gaspar EM, Duarte R, Santana JC. Volatile composition and antioxidant properties of clove products. *Biomedical Journal of Scientific and Technical Research*. 2018; 9(4):7270-6.

<http://dx.doi.org/10.26717/BJSTR.2018.09.001831>

31- Lee S, Najiah M, Wendy W, Nadirah M. Chemical composition and antimicrobial activity of the essential oil of *Syzygium aromaticum* flower bud (Clove) against fish systemic bacteria isolated from aquaculture sites. *Frontiers of Agriculture in China*. 2009; 3:332-6 <https://doi.org/10.1007/s11703-009-0052-8>

32- Wongsawan K, Chaisri W, Tangtrongsup S, Mekrirat R. Bactericidal effect of clove oil against multidrug-resistant *Streptococcus suis* isolated from human patients and slaughtered pigs. *Pathogens*. 2019 21;9(1):14 .

<https://doi.org/10.3390/pathogens9010014>

33- Alanazi AK, Alqasbi MH, Alrouji M, Kuriri FA, Almuhanna Y, Joseph B, Asad M. Antibacterial activity of *syzygium aromaticum* (clove) bud oil and its interaction with imipenem in controlling wound infections in rats caused by methicillin-resistant *Staphylococcus aureus*. *Molecules*. 2022 Dec 5;27(23):8551.

<https://doi.org/10.3390/molecules27238551>

34- Gürbüz M, Korkmaz Bİ. The anti-campylobacter activity of eugenol and its potential for poultry meat safety: A review. *Food Chemistry*. 2022 15; 394:133519.

<https://doi.org/10.1016/j.foodchem.2022.133519>

35- Jang HI, Rhee KJ, Eom YB. Antibacterial and antibiofilm effects of α -humulene against *Bacteroides fragilis*. *Canadian Journal of Microbiology*. 2020; 66(6):389-99.

<https://doi.org/10.1139/cjm-2020-0004>

36- Zhang X, Sun X, Wu J, Wu Y, Wang Y, Hu X, Wang X. Berberine damages the cell surface of methicillin-resistant *Staphylococcus aureus*. *Frontiers in microbiology*. 2020 28; 11:621.

<https://doi.org/10.3389/fmicb.2020.00621>

37- Mandal S, Mandal MD, Pal NK. Antibacterial potential of *Azadirachta indica* seed and *Bacopa monniera* leaf extracts against multidrug resistant *Salmonella enterica* serovar Typhi isolates. *Archives of Medical Science*. 2007 1; 3(1):14-8.

<https://www.termedia.pl/Original-paper-Antibacterial-potential-of-Azadirachta-indica-seed-and-Bacopa-monniera-leaf-extracts-against-multidrug-resistant-Salmonella-enterica-serovar-Typhi-isolates,19,7850,0,1.html>

38- Mandal S, DebMandal M, Saha K, Pal NK. In-vitro antibacterial activity of three Indian spices against methicillin-resistant *Staphylococcus aureus*. *Oman Medical Journal*. 2011; 26(5):319.

<https://doi.org/10.5001%2Fomj.2011.80>

How to Cite this Article:

Mays U. Hashim, M.S.Ibrahim O. Extraction and
Identification of the Main Components of Cloves
(*Syzygium Aromaticum* L.) Oil Extract and its
Antimicrobial Activity against MRSA strain.

JFacMedBagdad 66(2). Available

from: <https://iqjmc.uobaghdad.edu.iq/index.php/19JFacMedBaghdad36/article/view/2187>

استخراج وتحديد المكونات الرئيسية لمستخلص زيت نبات القرنفل (*Syzygium Aromaticum* L) وفعاليتها المضادة للميكروبات ضد جرثومة المكورات العنقودية الذهبية المقاومة للميثيسيلين.

ميس عدي هاشم¹, عروبة محمد سعيد ابراهيم²
1فرع الفلسفة والكيمياء والادوية, كلية الطب البيطري, جامعة القاسم الخضراء, بابل, العراق
2فرع الفلسفة والكيمياء والادوية, كلية الطب البيطري, جامعة بغداد, بغداد, العراق

الخلاصة:

خلفية: تُعرف بكتيريا المكورات العنقودية الذهبية المقاومة للميثيسيلين على نطاق واسع كعامل مسبب مهم مسؤول عن العدوى في جميع أنحاء العالم. وان واحدة من أكبر المشاكل في مجال الرعاية الصحية العالمية هي مقاومة المضادات الحيوية لهذه البكتيريا. حيث يعد استخدام الأدوية العشبية إحدى التقنيات الواعدة لمواجهة مقاومة البكتيريا للمضادات الحيوية.

الأهداف: دراسة التركيب الكيميائي واختبار النشاط المضاد للبكتيريا في المختبر لمستخلص زيت القرنفل ضد جرثومة المكورات العنقودية الذهبية المقاومة للميثيسيلين.

الطريقة: تم الحصول على مستخلص زيت القرنفل باستخدام التقطير المائي بواسطة جهاز Clevenger. بعد ذلك، تم إجراء التحليل الكيميائي النباتي عن طريق التحليل الكروماتوجرافي-مطياف الكتلة. تم إجراء فحص لمضادات الميكروبات لزيت نبات القرنفل في المختبر باستخدام طريقة انتشار القرص المزدوج، وطريقة التخفيف الدقيق، وطريقة منحنى وقت القتل. تم استخدام اختبار أقل فرق معنوي (ANOVA) LSD- (تحليل التباين) لإجراء مقارنة معنوية بين النتائج في هذه الدراسة.

النتائج: أظهرت هذه الدراسة ان نسبة استخلاص مستخلص القرنفل 50%. وقد أظهر التحليل الكيميائي النباتي وجود مكونات مختلفة وبشكل رئيسي هي بنسبة 28% Caryophyllen و 21.6% Humulene و 13.06% Eugenol. حيث تبين ان المكورات العنقودية الذهبية كانت حساسة لمستخلص القرنفل بتركيزات مختلفة علاوة على ذلك، وجد أن أقل تركيز مثبط (MIC) وأقل تركيز مبيد للجراثيم (MBC) كان 2.5 ميكروغرام / مل، 5.0 ميكروغرام / مل على التوالي. وقد حقق منحنى وقت القتل من xMICs2 و xMICs4 لمستخلص القرنفل أعلى تأثير قاتل للجراثيم مقارنة بالتركيز الأخرى. ($P \geq 0.05$)

الاستنتاجات: ظهر تحليل GC-MS لزيت القرنفل وجود 17 مكونًا متطابقًا في مستخلص القرنفل ويمكن أن تُعزى الخصائص المضادة للبكتيريا الجيدة التي أظهرها مستخلص زيت القرنفل ويعزى ذلك إلى المركبات الموجودة في الفينوليه مثل الكاريفيلين، الهيمولين، والوجينول.
الكلمات المفتاحية: المكورات العنقودية الذهبية المقاومة للميثيسيلين (MRSA)، الكاريفيلين، مضادات الميكروبات، مستخلص القرنفل، منحنى وقت القتل