

## Correlation between MDA Level and Chitotriosidase-1 Activity in Seminal Fluid of Iraqi Infertile Males

Ali S. Abdul Aziz<sup>1</sup> , Hedef Dh. El-Yassin<sup>\*1</sup> , Hussain K. Kadhem<sup>2</sup> 

<sup>1</sup>Department of Clinical Biochemistry, College of Medicine, University of Baghdad, Baghdad, Iraq.

<sup>2</sup>Department of Infertility, Al-Batool Teaching Hospital, Ministry of Health, Diyala, Iraq.



©2024 The Author(s). Published by College of Medicine, University of Baghdad. This open-access article is distributed under the terms of the Creative Commons Attribution 4.0 International License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

### Abstract

**Background:** Male infertility is a multifactorial condition influenced by various physiological and biochemical factors. Seminal fluid composition plays a crucial role in sperm function and fertilization potential. Chitotriosidase is a chitinase enzyme released by activated macrophages and is highly conserved and controlled. The notable chitinase in humans plays a significant role in the body's immunological response and is linked to inflammation, infection, tissue damage, and remodeling processes. On the other hand, malondialdehyde is a marker of lipid peroxidation, reflecting oxidative stress levels.

**Objective:** This study aimed to explore the correlation between malondialdehyde levels and Chitotriosidase-1 in seminal fluid in Iraqi infertile males.

**Methods:** Ninety males aged between twenty and forty-five were included in this cross-sectional study, all diagnosed with infertility by specialists at the infertility unit of Al-Batool Teaching Hospital between February 2022 and February 2023. The participants were categorized into three groups: the Normozoospermic Group (G1), the Asthenospermia Group (G2), and the Oligozoospermic Group (G3). Seminal malondialdehyde and Chitotriosidase-1 levels were measured by competitive Enzyme-linked immunosorbent assay.

**Results:** The study findings showed significantly higher levels of seminal fluid Chitotriosidase-1 found in the G2 group compared to the G3 and G1 groups. The seminal fluid malondialdehyde level for G1 was significantly lower than those for G2 and G3, which revealed a significant positive correlation between seminal fluid Chitotriosidase-1 activity and malondialdehyde levels ( $r = 0.37$ ,  $P < 0.05$ ) in the Asthenospermia Group.

**Conclusion:** There is a correlation between seminal fluid Chitotriosidase-1 activity and malondialdehyde level in the Asthenospermia Group. Novel diagnostic and therapeutic approaches for the treatment of male infertility may result from our growing understanding of the roles played by Chitotriosidase-1 and malondialdehyde in male reproductive health.

**Keywords:** Chitotriosidase-1; Male infertility; Malondialdehyde; Seminal plasma; Sperm quality.

Received: May, 2024

Revised: Aug. 2023

Accepted: Sept. 2024

Published: Dec.2024

### Introduction

Fertility is the ability of the individual to reproduce through normal sexual acts. Normal fertility requires the production of enough healthy sperm, a problem with this step causes infertility (1). Several studies were carried out on the association of infection and inflammation with male infertility, which revealed great variations in the prevalence of genital infection in different parts of the world (2). Involvement of chitotriosidase-1 (CHIT1) in macrophage activation and differentiation has consequences for other immune cell types. Although this may point to a role for CHIT1 in triggering an inflammatory response, data on the

enzyme's involvement in the inflammation that contributes to male infertility is still lacking (3). Chitotriosidase-1 is an enzyme found in various tissues and bodily fluids, including blood and seminal fluid. In humans, CHIT1 is primarily produced and secreted by macrophages, where it serves as part of the innate immune response against chitin-containing pathogens. Elevated levels of CHIT1 activity have been associated with various conditions, including lysosomal storage disorders and certain inflammatory diseases (4). Malondialdehyde (MDA) is the most well-studied by-product of polyunsaturated fatty acid peroxidation caused by oxidative damage (5). Seminal levels of reactive oxygen species and MDA also increase in tandem with these findings, during infection and tissue

\*Corresponding  
[hedefelyassin@uobaghdad.edu.iq](mailto:hedefelyassin@uobaghdad.edu.iq)

Author:

damage, semen contains high amounts of the same cytokines that play an important role in immune regulation for the male gonad (6). This study aimed to explore the correlation between malondialdehyde levels and Chitotriosidase-1 in seminal fluid in Iraqi infertile males.

## Patients and Methods:

### Study Population

Ninety males aged between twenty and forty-five were included in this cross-sectional study all were diagnosed with infertility by specialists at the infertility unit of Al-Batool Teaching Hospital / diyalah governorate between February 2022 and February 2023. Subjects were divided according to seminal fluid analysis into three groups: the Normozoospermia Group (G1), the Asthenospermia Group (G2), and the Oligozoospermia Group (G3).

Inclusion criteria (according to Seminal Fluid Analysis (SFA) (WHO,1999)

- ❖ Normozoospermic Group (G1): must have a normal sperm count, Sperm morphology Particiy (shape and structure) should meet standard criteria for normal sperm, Sperm motility (ability to move) should meet standard criteria for normal sperm.

- ❖ Asthenospermia Group (G2): must have reduced sperm motility, other parameters such as sperm count and morphology may still fall within the normal range.

- ❖ Oligozoospermic Group (G3): must have a low sperm count, Sperm morphology and motility may be normal.

Exclusion criteria

Systemic disease such as DM.

- 1- Auto immune disease such as SLE, RA, Hashimoto's thyroiditis.
- 2- Severe oligospermia 5 million / ejaculation.
- 3- Azoospermia
- 4- Patients with varicocele.
- 5- Patients with disorders in his wife reproductive system.
- 6- Patients with undescending testis and with testicular torsion.
- 7- Hypogonadism
- 8- Patients previously taken Antimicrobials, corticosteroid, and Antioxidants.

### Seminal fluid samples

All patients' semen samples were taken in sterile, clean cups, and placed in an incubator for 15-20 minutes to cause the semen to liquefy, throughout the course of three to four days of abstinence. The samples were then examined under a light microscope. Seminal plasma was obtained by centrifugation at 4000 rpm for 15 minutes were divided into two portions and kept until assay.

**Measurements of MDA and Chitinase1 (CHIT1) by enzyme-linked immunosorbent assay (ELISA)**

Seminal MDA and CHIT1 levels were measured by competitive Enzyme-linked immunosorbent assay (ELISA) according to the manufacturer (Cloud-Clone Corp/USA/Cat No. CEA597Ge, SEJ374Hu, SEA181Hu).

### Statistical Analysis:

The statistical analysis was conducted using Microsoft Excel for data input and preparation, which included organizing and cleaning the data for analysis, One-way ANOVA followed by multiple comparisons test was performed using GraphPad Prism version 19.5.1 for Windows, GraphPad Software, San Diego, California USA, and MedCalc® Statistical Software version 20.215 was used to calculate and examine the strength and direction of relationships between variables, particularly between seminal plasma Malondialdehyde (SF-MDA) and CHIT1 levels. The statistical significance level was set at 0.05 for most tests, indicating that results with a [P-value less than 0.05 were considered statistically significant.

### Results:

Age appeared to be uniformly distributed across the three groups with mean ages of  $29.67 \pm 1.043$ ,  $29.86 \pm 0.896$ , and  $29.96 \pm 0.752$  years, for G1, G2, and G3, respectively. The  $P=0.972$ , strongly suggests that the differences between these groups were not statistically significant, demonstrated more discernible differences among the groups as shown in figure (1).




Figure 1: Age of males in the study groups.

In table 1 and figure (2), the results showed, that the Seminal Fluid Volume (SF Vol) revealed mean values of  $2.550 \pm 0.1701$ ,  $2.083 \pm 0.1337$ , and  $2.033 \pm 0.1809$  ml for study groups, respectively. Although differences were observed, the effect size indicated by the  $R^2$  value of 0.07 and a borderline  $P=0.05$  suggested that these differences are not substantial.

For Sperm Count, however, a striking disparity among the groups was evident. The means were  $71.833 \pm 1.8489$ ,  $39.333 \pm 2.3700$ , and  $11.000 \pm 0.7350$  for

study groups, respectively. The  $R^2$  value of 0.87 and a  $P= 0.001$  indicated not only a statistically significant difference but also a large effect size, denoting a considerable variation in sperm count among these groups.


Motility categories, designated as A%, B%, C%, and D%, also showed substantive differences among the groups. For category A%, the mean values were  $21.833\pm 0.9123$ ,  $6.500\pm 1.0491$ , and  $2.000\pm 0.9160$  for study groups, respectively. Category B% displayed similar disparities with means of  $33.833\pm 0.5715$ ,  $18.000\pm 1.0057$ , and  $8.833\pm 1.6380$ . Both A% and B% had  $R^2$  values of 0.73 and  $P$ -values less than 0.001,

signifying statistical significance and substantial effect sizes. Category C%, while also statistically significant with  $P= 0.01$ , showed a modest  $R^2$  value of 0.10, implying a smaller effect size. Category D% exhibited an  $R^2$  value of 0.77 and a  $P$ -value less than 0.001, indicating significant differences with a large effect size. Finally, the percentages of morphologically normal and abnormal sperms were  $70.000\pm 0.8970$  and  $30.333\pm 0.8949$  for G1,  $40.833\pm 2.5380$  and  $59.167\pm 2.5380$  for G2, and  $23.833\pm 3.5934$  and  $76.167\pm 3.5934$  for G3, respectively. These differences were statistically significant with  $P= 0.001$  and  $R^2$  values of 0.65, highlighting a sizable effect size.

**Table 1 Semen profile for study groups**

SFA	Study groups			ANOVA	
	G1	G2	G3	$R^2$	P value
SF Vol ml	$2.550\pm 0.1701$	$2.083\pm 0.1337$	$2.033\pm 0.1809$	0.07	0.05
Sperm count	$71.833\pm 1.8489$	$39.333\pm 2.3700$	$11.000\pm 0.7350$	0.87	< 0.001
A%	$21.833\pm 0.9123$	$6.500\pm 1.0491$	$2.000\pm 0.9160$	0.73	< 0.001
B%	$33.833\pm 0.5715$	$18.000\pm 1.0057$	$8.833\pm 1.6380$	0.73	< 0.001
C%	$10.000\pm 0.0000$	$10.833\pm 0.3460$	$9.333\pm 0.4632$	0.10	0.01
D%	$34.333\pm 1.0095$	$64.667\pm 1.9613$	$79.833\pm 2.5312$	0.77	< 0.001
Normal%	$70.000\pm 0.8970$	$40.833\pm 2.5380$	$23.833\pm 3.5934$	0.65	< 0.001
Abnormal%	$30.333\pm 0.8949$	$59.167\pm 2.5380$	$76.167\pm 3.5934$	0.65	< 0.001

SFA=Seminal fluid analysis, SF vol=Seminal fluid volume, SE= Standard error of mean, and Motility categories=A%, B%, C%, and D%




**Figure 2: Means of Semen profile by the study groups.**

Seminal fluid plasma Ch1t1 levels also displayed pronounced disparities among the groups with mean concentrations of  $5.62\pm 0.19$ ,  $15.96\pm 0.29$ , and  $12.46\pm 0.18$  (ng/ml) for study groups, respectively. As seen Table 2 and Figure 3

**Table 2: The SF- Ch1t1 for study groups**

Parameter	Mean $\pm$ SE			P value
	G1	G2	G3	
SF_Ch1t1 (ng/ml)	$5.62\pm 0.19$	$15.96\pm 0.29$	$12.46\pm 0.18$	< .001




**Figure3: Means of Ch1t1 by study groups.**

The mean of SF-MDA for G1  $236.59 \pm 105.30$ (ng/ml), was significantly lower than for G2  $1904.08 \pm 877.85$ , similarly, compared to G3  $1,042.78 \pm 339.03$  with  $P=0.001$ . As seen Table 3, and Figure 4.

**Table 3: The SF-MDA for study groups**

Parameter	Group	n	Mean± SD	G1	G2	G3	Pr > F(Model)	Significant
SF-MDA ng/ml	G1	30	236.59± 105.30					
	G2	30	1904.08± 877.85	236.59 a	1904.08 c	1042.78 b	< 0.001	Yes
	G3	30	1042.78± 339.03					



**Figure 4: Comparison of SF-MDA between study groups.**

The statement highlights a significant and substantial difference in sperm count among the Normozoospermic Group (G1), Asthenospermic Group (G2), and Oligozoospermic Group (G3). This indicated that the observed variation in sperm count is unlikely to have occurred by chance alone.

The correlation between SF MDA and SF Ch1t1 is modestly positive ( $r=0.37$ ) and statistically significant ( $P$  value = 0.042) in G2 group. This suggested that there is a significant, albeit weak, positive association between SF MDA and SF Ch1t1 levels in the studied population. The modestly positive correlation between SF MDA and SF Ch1t1 may suggest a potential relationship between oxidative stress (MDA) and Ch1t1 activity in the seminal fluid.

**Table 4: Correlation between SF Ch1t1 and SF\_MDA in G2 group and G3**

Parameter	SF_Ch1t1 in G2		
	R	95.00% CI	p
SF_MDA	.37	[.01, .65]	.042
Parameter	SF_Ch1t1 in G3		
	R	95.00% CI	p
SF MDA	-.12	[-.46, .25]	.536

## Discussion:

The observed variation is substantial and not merely a result of the sample size, a considerable difference in sperm count among the groups has important clinical implications, sperm count is a crucial factor in male fertility, and significant variations may affect the likelihood of successful conception, the study may explore potential causes or factors contributing to the observed differences in sperm count (7).

Factors such as lifestyle, environmental exposures, and genetic predispositions could be the cause of such variations among the different sperm health conditions, there are several reasons for a significant difference in sperm count among the groups included in the study (8). These factors may result from complex interactions between genetic, environmental, and health-related elements, variations in levels of hormones involved in sperm production, such as testosterone, may contribute to this difference, differences in genetic makeup could affect sperm count, genetics play a crucial role in determining reproductive characteristics(9).

For seminal fluid volume, although there are differences, the  $P$ -value (Table 1;  $P \leq 0.05$ ) suggested that , there can be several reasons for differences in semen volume among different groups, including disruptions in hormone levels, particularly those involved in sperm production, can affect semen volume, imbalances in hormones like testosterone may influence semen production, variations in genes among individuals can contribute to differences in semen volume, genetic factors play a role in the physical characteristics of semen (10).

Exposure to specific environmental factors may influence natural sperm production and, consequently, semen volume. This could include exposure to harmful chemicals or elevated temperatures, as shown by (11). The provided data revealed significant differences in motility categories (A%, B%, C%, and D%) among the study groups, namely the Normozoospermic Group (G1), Asthenospermic Group (G2), and Oligozoospermic Group (G3).

The statistically significant differences in Category A% suggested that there is a significant variation in the percentage of sperms showing progressive motility among the different groups ( $P < 0.001$ ). This is a crucial parameter for male fertility, as sperm with progressive motility have a higher chance of reaching and fertilizing the egg (12).

Similar to Category A%, the significant differences ( $P < 0.001$ ) with a substantial effect size in Category B% indicated notable disparities in the percentage of sperms showing non-progressive motility. Non-progressive motility may still allow sperm to move but with less efficiency compared to progressive motility (13). Although statistically significant ( $P < 0.01$ ), the smaller effect size in Category C% suggested a less substantial impact compared to Categories A% and B%. This indicated that the percentage of sperms with local motility varies among the groups but to a lesser degree. Local motility may have limited functional relevance for fertilization compared to progressive motility (14). The significant differences ( $P < 0.001$ ) with a large effect size in Category D% highlighted considerable variations in the percentage of immotile sperm among the study groups. Immotile sperms have reduced or no movement, which can significantly impact fertility (15). The findings have clinical relevance, as sperm motility is a critical factor in male fertility, understanding the specific patterns of motility among different sperm health conditions can guide clinicians in diagnosing and addressing fertility issues in couples (16), as found by (17). With advancing age, there can be an effect on sperm motility, aging is often associated with a decline in overall body functions.

The provided information indicated significant differences in the percentages of morphologically normal and abnormal sperms among the three groups (G1, G2, and G3), these morphological differences may have implications for fertility and reproductive health in each group, this agreed with (18).

This finding can be important for the study's validity and interpretation, it suggested that the differences in sperm characteristics observed among the groups are more likely to be related to the specific condition (G1, G2, and G3) rather than being confounded by age or BMI differences (19).

As shown in Table (1) and Figure (4), seminal plasma MDA levels for G1 were significantly lower ( $P < 0.001$ ) than those for both G2 and G3.

Malondialdehyde (MDA) is a naturally occurring compound that serves as a marker for oxidative stress and lipid peroxidation in cells, it is a byproduct of the degradation of polyunsaturated fatty acids in cell membranes when cells are exposed to oxidative stress, such as from reactive oxygen species (ROS) (20), lipid peroxidation can occur, leading to the formation of MDA, this agrees with (21). Lower MDA levels in G1 might suggest reduced oxidative stress in the Normozoospermic Group (G1) compared to the Asthenospermic Group (G2) and Oligozoospermic Group (G3), oxidative stress is known to negatively impact sperm quality and function, high levels of ROS can lead to DNA damage, lipid peroxidation, and impaired sperm function. The observed difference in MDA levels among the groups may be associated with

the differences in sperm parameters, especially considering that G2 is associated with asthenospermia (reduced sperm motility) (22).

Elevated MDA levels are associated with various pathological conditions, including inflammation, cardiovascular diseases, neurodegenerative disorders, and reproductive health issues (23).

Studies often investigate MDA levels concerning male fertility. Elevated MDA levels in seminal plasma or sperm cells may be linked to reduced sperm motility, viability, and overall sperm function, this agreed with (24, 25).

The study demonstrated significant differences in CHIT1 concentrations in seminal plasma between the normozoospermic and oligozoospermic groups, which aligned with the findings of this study. Inflammation in the male reproductive system can cause oligozoospermia (26).

This suggests a potential link between chitin metabolism and oxidative stress in the male reproductive system. Furthermore, subgroup analysis based on semen parameters revealed varying correlations between CHIT1 and MDA levels among different fertility profiles.

Malondialdehyde is a marker of oxidative stress, and elevated levels of MDA in seminal fluid have been associated with decreased sperm quality and male infertility. Asthenospermia, a condition characterized by reduced sperm motility, can be influenced by oxidative stress (20)

Ch1t1, on the other hand, is a protein involved in sperm maturation and function. Studies have shown that alterations in Ch1t1 levels in seminal fluid may be associated with male infertility and impaired sperm function (3).

In the context of asthenospermia, it is possible that elevated levels of MDA and alterations in Ch1t1 levels in seminal fluid could contribute to the condition. Oxidative stress induced by high MDA levels may affect sperm motility, while changes in Ch1t1 levels could impact sperm maturation and function.

### Limitations:

There were some factors and reasons behind the limitations of the study: The small number of sample size that we were able to collect in the study, sample contamination was an obstacle to collecting a larger number of samples, Community customs in collecting semen samples restricted obtaining other samples, Some patients not providing their personal information to obtain the sample prevented obtaining more samples and information and the limited geographical area affected the generalization of the study to all patients in Iraq.



### Conclusion:

The observed correlation between seminal fluid CHIT1 activity and MDA levels underscores the interplay between chitin metabolism and oxidative stress in male fertility. Understanding the role of CHIT1 and MDA in male reproductive health could lead to novel diagnostic and therapeutic strategies for male infertility management.

### Authors' declaration:

We confirm that all the Figures and Tables in the manuscript belong to the current study. Besides, the Figures and images, which do not belong to the current study, have been given permission for re-publication attached to the manuscript. Authors sign on ethical consideration's approval-Ethical Clearance: The project was approved by the local ethical committee in (Department of Biochemistry) according to the code number (19634) on (20/ 5/ 2024).

**Funding:** None

**Conflicts of Interest:** None

### Authors' contributions:

Study conception & design: (Ali S. Abdul Aziz, Hedef Dhafir El-Yassin, HK Kadhem). Literature search: (Ali S. Abdul Aziz, Hedef Dhafir El-Yassin, HK Kadhem). Data acquisition: (Ali S. Abdul Aziz, Hedef Dhafir El-Yassin, HK Kadhem). Data analysis & interpretation: (Ali S. Abdul Aziz, Hedef Dhafir El-Yassin, HK Kadhem). Manuscript preparation: (Ali S. Abdul Aziz, Hedef Dhafir El-Yassin, HK Kadhem). Manuscript editing & review: (Ali S. Abdul Aziz, Hedef Dhafir El-Yassin, HK Kadhem).

### References

- 1- Al-Khateeb S, Hussein SM, Dahy AAAI. Evaluation of inhibin-B hormone, FSH, and Testosterone in serum of infertile men. *J Fac Med Baghdad* . 2016 Jul 3 ;58(2):180–2.  
<https://doi.org/10.32007/jfacmedbagdad.5822377>
- 2- Lazem AA, Al-Kaseer E, Al-Diwan JK, Al- Hadithi TS. Effect of infection on semen parameters in a sample of Iraqi infertile males. *J Fac Med Baghdad*. 2010;52(3):274–6.  
<https://doi.org/10.32007/jfacmedbagdad.523973>
- 3- Kratz EM, Zurawska-Plaksej E, Solkiewicz K, Kokot I, Faundez R, Piwowar A. Investigation of seminal plasma chitotriosidase-1 and leukocyte elastase as potential markers for “silent” inflammation of the reproductive tract of the infertile male - a pilot study. *J Physiol Pharmacol* . 2020;71(3):1–7.  
<https://doi.org/10.26402/jpp.2020.3.04>
- 4- Di Francesco AM, Verrecchia E, Manna S, Urbani A, Manna R. The chitinases as biomarkers in immune-mediate diseases. *Clin Chem Lab Med* . 2022 Jul 1;61(8):1363–81.  
<https://doi.org/10.1515/cclm-2022-0767>

- 5- Toto A, Wild P, Graille M, Turcu V, Crézé C, Hemmendinger M, et al. Urinary Malondialdehyde (MDA) Concentrations in the General Population-A Systematic Literature Review and Meta-Analysis. *Toxics* . 2022 Apr 1;10(4).  
<https://doi.org/10.3390%2Ftoxics10040160>
- 6- Morselli S, Sebastianelli A, Liaci A, Zaccaro C, Pecoraro A, Nicoletti R, et al. Male reproductive system inflammation after healing from coronavirus disease 2019. *Andrology* . 2022 Sep 1;10(6):1030–7.  
<https://doi.org/10.1111/andr.13138>
- 7- Al-Darawsha, T. Z., Dayioglu, N., Al-Azzawi, B. R., & Irez, T. Study a relationship between age, body mass index, and sperm parameters with sperm DNA fragmentation levels in Iraqi infertile patients. *Al-Ameed Journal for Medical Research and Health Sciences*, 2023;1(2), 3.  
<http://dx.doi.org/10.61631/3005-3188.1007>
- 8- Salman, F. S., Al-Qadhi, H. I., & Al Kareem, B. A. N-acetyl cysteine's effect on semen parameters in a sample of Iraqi men with oligoasthenoteratozoospermia. *Journal of the Faculty of Medicine Baghdad*,2022; 64(3), 170-174.  
<https://doi.org/10.32007/jfacmedbagdad.6431938>
- 9- Fink, J., & Horie, S. *The Multiple Health Benefits of Testosterone*. Cambridge Scholars Publishing,2022.  
<https://www.cambridgescholars.com/product/978-1-5275-7637->
- 10- Elfateh, F., Wang, R., Zhang, Z., Jiang, Y., Chen, S., & Liu, R. Influence of genetic abnormalities on semen quality and male fertility: A four-year prospective study. *Iranian Journal of Reproductive Medicine*,2014; 12(2), 95.  
<http://www.ncbi.nlm.nih.gov/pmc/articles/pmc4009560/>
- 11- Murgia, F., Corda, V., Serrenti, M., Usai, V., Santoru, M. L., Hurt, K. J. & Monni, G. Seminal fluid metabolomic markers of oligozoospermic infertility in humans. *Metabolites*,2020; 10(2), 64.  
<https://doi.org/10.3390%2Fmetabo10020064>
- 12- Haugen, T. B., Witczak, O., Hicks, S. A., Björndahl, L., Andersen, J. M., & Riegler, M. A. Sperm motility assessed by deep convolutional neural networks into WHO categories. *Scientific Reports*,2023; 13(1), 14777. <https://doi.org/10.1038/s41598-023-41871-2>
- 13- Lu, H., Xu, D., Zhao, L., Ruan, H., Wang, A., Hu, J. & Lu, W. Exploring the regulatory role of Linc00893 in asthenozoospermia: Insights into sperm motility and SSC viability. *Molecular Medicine Reports*,2024; 29(2), 1-12. <https://doi.org/10.3892/mmr.2023.13143>
- 14- Diyasa IGSM, Saputra WSJ, Gunawan AAN, Herawati D, Munir S, Humairah S. Abnormality Determination of Spermatozoa Motility Using Gaussian Mixture Model and Matching-based Algorithm. *J Robot Control [Internet]*. 2024 Jan 12 [cited 2024 Sep 3];5(1):103–16. Available from:

<https://journal.umy.ac.id/index.php/jrc/article/view/20686>

15- Dcunha R, Hussein RS, Ananda H, Kumari S, Adiga SK, Kannan N, Zhao Y, Kalthur G. Current Insights and Latest Updates in Sperm Motility and Associated Applications in Assisted Reproduction. *Reprod Sci*. 2022 Jan;29(1):7-25.

<https://doi.org/10.1007%2Fs43032-020-00408-y>

16- Schlegel, P. N., Sigman, M., Collura, B., De Jonge, C. J., Eisenberg, M. L., Lamb, D. J., ... & Zini, A. Diagnosis and treatment of infertility in men: AUA/ASRM guideline part I. *The Journal of urology*, 2021; 205(1), 36-43.

<https://doi.org/10.1097/ju.0000000000001521>

17- Pereira, P. P. D. S., Da Mata, F. A., Figueiredo, A. C. G., de Andrade, K. R. C., & Pereira, M. G. Maternal active smoking during pregnancy and low birth weight in the Americas: a systematic review and meta-analysis. *Nicotine & tobacco research*, 2017; 19(5), 497-505.

<https://doi.org/10.1093/ntr/ntw228>

18- Zanetti, B. F., Braga, D. P. A. F., Provenza, R. R., Figueira, R. C. S., Iaconelli Jr, A., & Borges Jr, E. Sperm morphological normality under high magnification is correlated to male infertility and predicts embryo development. *Andrology*, 2018; 6(3), 420-427. <https://doi.org/10.1111/andr.12473>

19- Farah Saad Hasan. Seminal Fluid Abnormality among Infertile Males: A Two-Center Based Study in Baghdad. *Iraqi Postgrad Med J* [Internet]. 2020 [cited 2024 Aug 30];19(2). Available from:

<https://www.iasj.net/iasj/article/186077>

20- Hassan ZM, Hamdi RA, Bassam EN Al. Evaluation of the Role of Serum Malondialdehyde in the Pathogenesis of Diabetic Retinopathy. *J Fac Med Baghdad* [Internet]. 2022 Oct 17 [cited 2024 Aug

30];64(3):195–8. Available from: <http://dx.doi.org/10.32007/jfacmedbagdad.6431957>

21- Kodali ST, Kauffman P, Kotha SR, Yenigalla A, Veeraraghavan R, Pannu SR, et al. Oxidative Lipidomics: Analysis of Oxidized Lipids and Lipid Peroxidation in Biological Systems with Relevance to Health and Disease. 2020 Aug 9;61–92. Available from:

<https://www.ncbi.nlm.nih.gov/books/NBK566435/>

22- Jasem KM, Alnasrawi TH, Shiblawi HH, Wahid HHA, Al-Saadi NH. Investigation of malondialdehyde and some elements in young infertile males. *Res J Pharm Technol*. 2021 Oct 1;14(10):5418–22. <http://dx.doi.org/10.52711/0974-360X.2021.00944>

23- Singh A, Kukreti R, Saso L, Kukreti S. Oxidative Stress: A Key Modulator in Neurodegenerative Diseases. *Molecules* [Internet]. 2019 Apr 22 [cited 2024 Aug 30];24(8). Available from:

<https://pubmed.ncbi.nlm.nih.gov/31013638/>

24- Kurkowska W, Bogacz A, Janiszewska M, Gabrys E, Tiszler M, Bellanti F, et al. Oxidative Stress is Associated with Reduced Sperm Motility in Normal Semen. *Am J Mens Health* [Internet]. 2020 Sep 1 [cited 2024 Aug 30];14(5). Available from:

<https://pubmed.ncbi.nlm.nih.gov/32938274/>

25- Bergsma AT, Li HT, Eliveld J, Bulthuis MLC, Hoek A, van Goor H, et al. Local and Systemic Oxidative Stress Biomarkers for Male Infertility: The ORION Study. *Antioxidants* [Internet]. 2022 Jun 1 [cited 2024 Aug 30];11(6). Available from:

<https://pmc/articles/PMC9220279/>

26- Hasan H, Bhushan S, Fijak M, Meinhardt A. Mechanism of Inflammatory Associated Impairment of Sperm Function, Spermatogenesis and Steroidogenesis. *Front Endocrinol (Lausanne)*. 2022 Apr 28;13:897029. <https://doi.org/10.3389/fendo.2022.897029>

#### How to Cite this Article

Abdul Aziz AS, Elyaseen HD, Kadhem HK. Estimation of the Seminal Fluid Chitotriosidase-1 and Malondialdehyde level in infertile Iraqi male with silent inflammation. *J Fac Med Baghdad* [Internet]. Available from: <https://iqjmc.uobaghdad.edu.iq/index.php/19JFacMedBaghdad36/article/view/2395>

## العلاقة بين مستوى كيتوتريوسايديز-1 والمالونديالدهيد في السائل المنوي للذكور العراقيين العقيمين

علي صكبان عبد العزيز<sup>1</sup>، هدف ظافر الياسين<sup>1</sup>، حسين خليفة كاظم<sup>2</sup>  
<sup>1</sup>فرع الكيمياء الحياتية، كلية الطب، جامعة بغداد، بغداد، العراق.  
<sup>2</sup>مستشفى البتول التعليمي، دائرة صحة ديالى، وزارة الصحة، ديالى، العراق.

### الخلاصة

**خلفية البحث:** العقم عند الرجال هو حالة متعددة العوامل تتأثر بعوامل فسيولوجية وكيميائية حيوية مختلفة. يلعب تكوين السائل المنوي دوراً حاسماً في وظيفة الحيوانات المنوية وإمكانية الإخصاب. كيتوتريوسايديز-1 هو إنزيم يشارك في استقلاب الكيتين، في حين أن المالونديالدهيد (هو علامة على بيروكسيد الدهون، مما يعكس مستويات الإجهاد التأكسدي). تهدف هذه الدراسة إلى استكشاف العلاقة بين مستويات المالونديالدهيد و كيتوتريوسايديز ( في السائل المنوي عند الذكور العراقيين المصابين بالعقم والذين يعانون من التهاب صامت).

**طرق العمل:** تم تضمين تسعين رجلاً تتراوح أعمارهم بين عشرين وخمسة وأربعين عاماً في هذه الدراسة المقطعية، وتم تشخيص جميعهم بالعقم من قبل متخصصين في وحدة العقم في مستشفى البتول التعليمي بين فبراير 2022 وفبراير 2023. وتم تصنيف المشاركين إلى ثلاث مجموعات: مجموعة طبيعية النطاف (G1)، ومجموعة وهن النطاف (G2)، ومجموعة قليلة النطاف (G3).

**النتائج:** النتائج التي توصلنا إليها، كانت هناك مستويات أعلى بكثير من كيتوتريوسايديز في السائل المنوي الموجودة في المجموعة G2 مقارنة بمجموعتي G3 و G1. كانت مستويات المالونديالدهيد للسائل المنوي لـ G1 أقل بكثير من تلك الخاصة بـ G2 و G3. كشفت عن وجود علاقة إيجابية مهمة بين نشاط السائل المنوي كيتوتريوسايديز ومستويات المالونديالدهيد ( $p < 0.05$ ,  $r = 0.37$ ) في مجموعة قليلة النطاف.

**الاستنتاجات:** وجود علاقة إيجابية معنوية بين نشاط السائل المنوي بمستويات المالونديالدهيد ومستويات كيتوتريوسايديز في مجموعة وهن النطاف.

**الكلمات المفتاحية:** السائل المنوي البشري، كيتوتريوسايديز-1، الالتهاب الصامت، المالونديالدهيد، بيروكسيد الدهون